arXiv:1504.04448v2 [math.RA] 20 Jul 2015

ON n-CUBIC PYRAMID ALGEBRAS

Abstract. In this paper we study a class of algebras having n-dimensional pyramid shaped quiver with n-cubic cells, which we called n-cubic pyramid algebras. This class of algebras includes the quadratic dual of the basic n-Auslander absolutely n-complete algebras introduced by Iyama. We show that the projective resolution of the simples of n-cubic pyramid algebras can be characterized by n-cuboids, and prove that they are periodic. So these algebras are almost Koszul and $(n-1)$-translation algebras. We also recover Iyama's cone construction for n-Auslander absolutely n-complete algebras using n-cubic pyramid algebras and the theory of n-translation algebras.

1. Introduction

Quivers, especially translation quivers, are very important in the representation theory of algebras [1, 4, 21]. There are many algebras, such as Auslander algebras, preprojective algebras, related to translation quivers [4, 9, 6]. They are widely used in many mathematical fields, such as Cohen-Maucauley modules, cluster algebras, Calabi-Yau algebras and categories, non-commutative algebraic geometry and mathematical physics [3, 5, 19, 20, 10, 8, 2]. Recently, Iyama has developed the higher representation theory [16, 17, 18], where a class of higher translation quivers also plays an important role. In [18], he characterizes a class of higher representation-finite algebras, n-Auslander absolutely n-complete algebras. We observed that the quivers of such algebras are n-dimensional pyramid shaped quivers with n-cubic cells. We study in this paper a class of algebras, call n-cubic pyramid algebras, defined on such quivers, which includes quadratic dual of n-Auslander absolutely n-complete algebras. We draw the quiver in a different way to show that the arrows in such quiver in fact point to n-directions in a n-dimensional space.

Almost Koszul rings are introduced by Brenner, Butler and King in [7]. One of their main results in the paper is the periodicity of the trivial extensions of representation finite hereditary algebras of bipartite oriented quiver. In fact, they prove that such algebras are almost Koszul of type $(3, h-1)$ for the Coxeter number h of the quiver of the algebra. Our aim is to generalize a modified version of this result to n-cubic pyramid algebras. We proved that a stable n-cubic pyramid algebra of height $m \geq 3$, that is, a twisted trivial extension of a corresponding n-cubic pyramid algebra, is almost Koszul of type $(n+1, m-1)$. Our proof is based on the combinatoric characterization of the projective resolutions of the simple modules of the stable n-cubic pyramid algebras, using the integral points on an $(n+1)$-cuboid defined on the corresponding vertex in the quiver. Observe that the class of trivial extensions of hereditary algebras of bipartite oriented quiver is the same as the trivial extensions of the quadratic dual of the hereditary algebras. Our result can be regarded as a higher representation theory version of the result of Brenner, Butler and King [7] in the case A_m.

This work is partly supported by Natural Science Foundation of China #11271119, and by Provincial Innovation Foundation For Postgraduate of Hunan #CX2014B189.

In [11], one of the authors introduces translation algebras as algebras with translation quivers as their quivers and the translation corresponds to an operation related to the Nakayama functor. Such algebras includes the quadratic dual of Auslander algebras, preprojective algebras. In [15], n-translation quivers and n-translation algebras are introduced. The classical construction of $\mathbb{Z}Q$ is generalized to construct an $(n+1)$-translation algebra from an n-translation algebra with admissible n-translation quiver, using trivial extension followed by a smash product with $\mathbb{Z}$. In this paper, we use this method and a truncation called cuboid truncation to construct $(n+1)$-cubic pyramid algebra from n-cubic pyramid algebra. By taken quadratic dual of a special case, we recover Iyama's cone construction of absolutely $(n+1)$-complete algebra $T_m^{(n+1)}(k)$ from an absolutely n-complete algebra $T_m^{(n)}(k)$ in [18].

The paper is organized as follow. In Section 2 we introduce pyramid shaped n-cubic quivers and corresponding algebras. We also introduce the related concepts like stable pyramid shaped n-cubic quiver for the description of the twisted trivial extensions, and $(p+1)$-cuboid for characterizing the minimal projective resolutions of the simples. Section 3 is devoted to characterize the linear part of this projective resolution of the simples of the stable n-cubic pyramid algebras using $(n+1)$-cuboid. As a corollary, we prove that the twisted trivial extensions of n-cubic pyramid algebra is $(n+1, q)$-Koszul. In Section 4, we show how to truncate an $(n+1)$-cubic pyramid algebra from the smash product of the twisted trivial extension of an n-cubic pyramid algebra. We also prove that $(n+1)$-cubic pyramid algebra is an n-translation algebra. In the last section, we study the higher almost split sequence in the category of the finitely generated projective modules for an n-cubic pyramid algebra and its quadratic dual. We also show how n-cubic pyramid algebras are used to recover Iyama's construction of absolutely n-complete algebras $T_m^{(n)}(k)$.

2. Pyramid Shaped n-cubic Quivers and Related Algebras

Let k be a field. In this paper, algebra is assumed to be a graded quotient of a path algebra of a locally finite quiver Q over k, that is, $\Lambda = kQ/(\rho) = \Lambda_0 + \Lambda_1 + \cdots$, with Λ_0 a direct sum of (possibly, infinite) copies of k and Λ generated by Λ_1 over Λ_0, with relation set ρ (see [15]). We assume the vertex set of Q is Q_0 and the arrow set is Q_1. So we have a complete set of idempotents $\{e_i | i \in Q_0\}$ such that $\Lambda_0 = \bigoplus_{i \in Q_0} \Lambda_0 e_i$ and the number of arrows from i to j is exactly $\dim_k e_j \Lambda_1 e_i$ for any $i, j \in Q_0$.

To introduce pyramid shaped n-cubic quiver, we first make some convention on the non-negative integral vectors.

Given a positive integer p, let $\mathbf{e}_0 = 0$, $\mathbf{e}^{(p)} = (1, \ldots, 1)$ be the p-dimensional vector with all the components 1 and let $\mathbf{e}_t^{(p)} = (0, \cdots, 0, 1, 0, \cdots, 0)$ be the p-dimensional vector with tth component 1 and all the other components zero. Write $\mathbf{e}_J^{(p)} = \mathbf{e}_{t_1,\ldots,t_s}^{(p)} = \sum_{h=1}^{s} \mathbf{e}_{t_s}^{(p)}$ for a subset $J = \{t_1, \ldots, t_s\}$ of $\{1, \ldots, p\}$, set $\bar{\mathbf{e}}_t^{(p)} = \mathbf{e}_{\{t,t+1\}}^{(p)}$. Conventionally, we write them as $\mathbf{e}, \mathbf{e}_t, \mathbf{e}_J$ and $\bar{\mathbf{e}}_t$ when no confusion will occur.

For a vector denoted by the boldface letter such like $\mathbf{a}$ write a_s for its sth component, so $\mathbf{a} = (a_1, \ldots, a_p)$. For a vector $\mathbf{i} = (i_1, \ldots, i_p) \in \mathbb{Z}^p$, write $\mathbf{i}(1) = \mathbf{i} + \mathbf{e}_1$, $\mathbf{i}(t) = \mathbf{i} - \mathbf{e}_{t-1} + \mathbf{e}_t$ for $2 \le t \le p$, and write $\mathbf{i}(t_1, \ldots, t_s) = \mathbf{i}(t_1) \ldots (t_s)$.

Similarly, write $(1)\mathbf{i} = \mathbf{i} - \mathbf{e}_1$, $(t)\mathbf{i} = \mathbf{i} + \mathbf{e}_{t-1} - \mathbf{e}_t$ for $2 \leq t \leq p$, and write $(t_1, \ldots, t_s)\mathbf{i} = (t_s) \ldots (t_1)\mathbf{i}$. Clearly $(t)\mathbf{i}(t) = \mathbf{i}$.

Let $\mathbb{Z}^+$ be the set of non-negative integers. For $\mathbf{a} = (a_1, \cdots, a_p) \in \mathbb{Z}^{+p}$, write $|\mathbf{a}| = \sum_{t=1}^{p} a_t$. Let $\mathbb{Z}^{+p}_l = \{\mathbf{a} = (a_1, \cdots, a_p) \in \mathbb{Z}^{+p} | |\mathbf{a}| = l\}$. We need the following subsets of $\mathbb{Z}^{+p}$. The vertex set

$$U^{(p)} = \{\mathbf{u} = (u_1, u_2, \cdots, u_p) | 0 \leq u_t \leq 1\} \subset \mathbb{Z}^{+p},$$

of the unit p-cube and the vertex sets

$$U^{(p)}_l = \{\mathbf{u} = (u_1, u_2, \cdots, u_p) | |\mathbf{u}| = l\},$$

of l-nets inside the unit p-cube, for $0 \leq l \leq p$.

Fix $m \geq 3$ and $\mathbf{i} = (i_1 \ldots, i_p) \in \mathbb{Z}^{+p}$ satisfying

$$1 \leq i_t \text{ and } \sum_{t=1}^{s} i_t \leq m + s - 1 \text{ for } s = 1, \ldots, p. \tag{1}$$

Set

$$b_1(\mathbf{i}) = m + p - 1 - |\mathbf{i}|, \quad b_t(\mathbf{i}) = i_{t-1} - 1, \quad 2 \leq t \leq p + 1,$$

write $b_t = b_t(\mathbf{i})$ and $\mathbf{b} = \mathbf{b}(\mathbf{i}) = (b_1, \ldots, b_{p+1})$. Let

$$C^{(p)}(\mathbf{i}) = \{(a_1, a_2, \cdots, a_{p+1}) | 0 \leq a_t \leq b_t(\mathbf{i}), 1 \leq t \leq p + 1\} \subset \mathbb{Z}^{+p+1},$$

this is the set of integral vertices of a $(p + 1)$-cuboid with sides of length b_t for $1 \leq t \leq p + 1$. We call it the $(p + 1)$-*cuboid associated to the vertex* $\mathbf{i}$. The vertex $C^{(p)}_0(\mathbf{i}) = (0, \ldots, 0)$ is called the initial vertex of the $C^{(p)}(\mathbf{i})$. Let

$$C^{(p)}_l(\mathbf{i}) = \{\mathbf{a} = (a_1, \cdots, a_{p+1}) \in C^{(p)}(\mathbf{i}) | \, |\mathbf{a}| = l\},$$

for $0 \leq l \leq m - 1$. $C^{(p)}_l(\mathbf{i})$ is a net inside the $(p + 1)$-cuboid formed by integral vertices which can be reached from the initial vertex in l steps. We have that $C^{(p)}_{m-1}(\mathbf{i}) = \{\mathbf{b} = (b_1, \ldots, b_{p+1})\}$, and $C^{(p)}_l(\mathbf{i}) = \emptyset$ for $l \geq m$.

For $\mathbf{i} = (i_1, \ldots, i_p) \in \mathbb{Z}^{+p}$ satisfying (1), for each $\mathbf{a} = (a_1, \cdots, a_{p+1}) \in \mathbb{Z}^{+p+1}$, define

$$\mathbf{v}^{\mathbf{i}}(\mathbf{a}) = (i_1 + a_1 - a_2, i_2 + a_2 - a_3, \cdots, i_{p-1} + a_{p-1} - a_p, i_p + a_p - a_{p+1}),$$

For $\mathbf{i} = (i_1, \ldots, i_p, i_{p+1}) = (\mathbf{i}', i_{p+1}) \in \mathbb{Z}^{+p+1}$ with $\mathbf{i}' = (i_1, \ldots, i_p)$ satisfying (1), write $C^{(p)}(\mathbf{i}) = C^{(p)}(\mathbf{i}')$. For each $\mathbf{a} = (a_1, \cdots, a_{p+1}) \in \mathbb{Z}^{+p+1}$, define

$$\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{a}) = (i_1 + a_1 - a_2, i_2 + a_2 - a_3, \cdots, i_p + a_p - a_{p+1}, i_{p+1} + a_{p+1}).$$

From now on, we fix integers $n \geq 1$ and $m \geq 3$. Define *n-cubic pyramid quiver* $Q(n)$ of height m as the quiver with

$$\begin{array}{ll} \text{vertex set:} & Q(n)_0 = \{\mathbf{i} = (i_1, \ldots, i_n) | 1 \leq i_t, \sum_{t=1}^{s} i_t \leq m + s - 1, 1 \leq s \leq n\} \\ \text{arrow set:} & Q(n)_1 = \{\gamma^{(t)}_{\mathbf{i}} : \mathbf{i} \longrightarrow \mathbf{i}(t) | 1 \leq t \leq n, \mathbf{i}, \mathbf{i}(t) \in Q(n)_0\}. \end{array} \tag{2}$$

Call $\gamma^{(t)}_{\mathbf{i}}$ an *arrow of type* t starting at $\mathbf{i}$. In the case of $n = 3$, these quivers look like a pyramid of side length of m built up with cubes:



These quivers are one of those defined in [18] inductively to describe the n-Auslander absolutely n-complete algebras. It is shown in [14] that such quivers are truncation of the McKay quivers of some finite Abelian subgroups of $GL_n(\mathbb{C})$.

For $\mathbf{i} = (i_1, \dots, i_n) \in Q(n)_0$, call the integral vector $\mathbf{a} \in \mathbb{Z}^{+n+1}$ an $\mathbf{i}$-*quiver vertex* if $\mathbf{v}^{\mathbf{i}}(\mathbf{a}) \in Q(n)_0$. Clearly, $\mathbf{v}^{\mathbf{i}}(\mathbf{a})$ is in $Q(n)_0$ if and only if $i_t > a_{t+1} - a_t$ and $a_{s+1} - a_1 \leq m + s - 1 - \sum\limits_{t=1}^{s} i_t$ for $1 \leq s \leq n$.

Let $\mathbf{i}$ be a vertex in $Q(n)$. The *n-cubic cell at* $\mathbf{i}$ *in* $Q(n)$ is the full subquiver $H^{\mathbf{i}}$ of $Q(n)$ with the vertex set the set of the $\mathbf{i}$-quiver vertices in $\{\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{a}) \in Q(n)_0 | \mathbf{a} \in U^{(n)}\}$. The vertex $\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{e})$ is called *the end vertex* of $H^{\mathbf{i}}$ and $H^{\mathbf{i}}$ is called *complete* if $\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{e})$ is a vertex of $H^{\mathbf{i}}$. $H^{\mathbf{i}}$ can be regarded as formed by certain vertices and the edges in the unit n-cube directed from 0 to $\mathbf{e}$. Define *n-hammock* $H^{\mathbf{i}}_C$ *at* $\mathbf{i}$ as the full subquiver of $Q(n)$ with the vertex set the set of the $\mathbf{i}$-quiver vertices in $\{\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{a}) \in Q(n)_0 | \mathbf{a} \in C^{(n-1)}(\mathbf{i})\}$. The vertex $\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{b}(\mathbf{i}))$ is called *the end vertex* of $H^{\mathbf{i}}_C$ and $H^{\mathbf{i}}_C$ is called *complete* if $\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{b}(\mathbf{i}))$ is a vertex of $H^{\mathbf{i}}_C$.

Let

$$\mathbf{g} = (g_1 = 1, g_2, \dots, g_n) \tag{3}$$

be a sequence of linear maps on $kQ(n)$ such that the restriction on the vertex set of g_s is only defined on the vertices with s component $i_s > 1$, in this case $g_s(\mathbf{i}) = \mathbf{i} - \mathbf{e}_s$, or, $g_s(e_{\mathbf{i}}) = e_{\mathbf{i}-\mathbf{e}_s}$, and $g_s(e_{\mathbf{i}}) = 0$ if $i_s = 1$. Let kQ_1 be the subspace of $kQ(n)$ spanned by the arrows of $Q(n)$, g_s restricts to a bijective linear map on $e_{\mathbf{i}} kQ_1 e_{\mathbf{j}}$ if $i_s \neq 1 \neq j_s$. Since there is at most one arrow between each pair of vertices, we have that $g_s(\gamma^{(t)}_{\mathbf{i}}) = d_{s,t,\mathbf{i}} \gamma^{(t)}_{g_s(\mathbf{i})}$ for some $0 \neq d_{s,t,\mathbf{i}} \in k$ when $i_s > 1$ and $t < s$.

Define a relation set:

$$\begin{aligned} \rho^{\mathbf{g}}(n) \quad = \quad & \{d_{s,t,\mathbf{i}} \gamma^{(t)}_{\mathbf{i}(s)} \gamma^{(s)}_{\mathbf{i}} - \gamma^{(s)}_{\mathbf{i}(t)} \gamma^{(t)}_{\mathbf{i}} | \mathbf{i}, \mathbf{i}(t), \mathbf{i}(s), \mathbf{i}(t)(s) \in Q(n)_0, 1 \leq t < s \leq n\} \\ & \cup \{\gamma^{(t)}_{\mathbf{i}(t)} \gamma^{(t)}_{\mathbf{i}} | \mathbf{i}, \mathbf{i}(t) \in Q(n)_0, 1 \leq t \leq n\}. \end{aligned} \tag{4}$$

The relations are of two kind, one is zero relation consisting of arrows of same type, and the other is commutative relation consisting of arrows of two different types.

Let $\Lambda^{\mathbf{g}}(n)$ be the algebra with bound quiver $(Q(n), \rho^{\mathbf{g}}(n))$, and we call it an *$n$-cubic pyramid algebra.* 1-cubic pyramid algebras are the quadratic duals of the hereditary algebras of quivers of type A_m with linear orientation.

A path p of a bound quiver (Q, ρ) is called a bound path if its image in $kQ/(\rho)$ is nonzero. Recall that a bound quiver (Q, ρ) is called an *n-translation quiver* [15] if there is a bijective map $\tau : Q_0 \setminus \mathcal{P} \longrightarrow Q_0 \setminus \mathcal{I}$, called the *$n$-translation* of Q, for two subsets $\mathcal{P}$ and $\mathcal{I}$ of Q_0, whose elements are called *projective vertices* and respectively *injective vertices*, satisfying the following conditions:

1. Any maximal bound path is of length $n+1$ from τi in $Q_0 \setminus \mathcal{I}$ to i, for some vertex i in $Q_0 \setminus \mathcal{P}$.
2. Two bound paths of length $n+1$ from τi to i are linearly dependent, for any $i \in Q_0 \setminus \mathcal{P}$.
3. For each $i \in Q_0 \setminus \mathcal{P}$ and $j \in Q_0$, any bound element u which is linear combination of paths of the same length $t \le n+1$ from j to i, there is a path q of length $n+1-t$ from τi to j such that uq is a bound element.
4. For each $i \in Q_0 \setminus \mathcal{I}$ and $j \in Q_0$, any bound element u which is linear combination of paths of the same length $t \le n+1$ from i to j, there is a path p of length $n+1-t$ from j to $\tau^{-1} i$ such that pu is a bound element.

An algebra Λ with bound quiver an n-translation quiver (Q, ρ) is called an *n-translation algebra* if there is an $q \in \mathbb{N} \cup \{\infty\}$ such that Λ is $(n+1, q)$-Koszul in the following sense

(1) $\Lambda_t = 0$ for $t > n+1$, and

(2) for each $i \in Q_0$, let

$$P^q(i) \xrightarrow{f_q} \cdots \longrightarrow P^1(i) \xrightarrow{f_1} P^0(i) \xrightarrow{f_0} S(i) = \Lambda_0 e_i \longrightarrow 0 \tag{5}$$

be the first $q+1$ terms in a minimal projective resolution of the simple $S(i) \simeq \Lambda_0 e_i$, then $P^t(i)$ is generated by its component of degree t for $0 \le t \le q$ and the kernel of f_q is concentrated in degree $n+1+q$.

Theorem 2.1. *$(Q(n), \rho^{\mathbf{g}}(n))$ is an admissible $(n-1)$-translation quiver with $(n-1)$-translation defined by $\tau : \mathbf{i} \longrightarrow \mathbf{i} - \mathbf{e}_n$ if $i_n > 1$.*

$\Lambda^{\mathbf{g}}(n)$ is an $(n-1)$-translation algebra with admissible $(n-1)$-translation quiver.

The theorem will be proven in the last section inductively (see Theorem 5.2).

Choosing $\tilde{\mathbf{g}} = (\tilde{g}_1, \ldots, \tilde{g}_n)$ such that $\tilde{g}_s(\gamma_{\mathbf{i}}^{(t)}) = -\gamma_{\tilde{g}_s(\mathbf{i})}^{(t)}$ for $t < s$ and for $\mathbf{i} = (i_1, \ldots, i_n)$, when all the arrows are in $Q(n)$, the following proposition follows directly.

Proposition 2.2. *The quadratic dual $\Lambda^{\tilde{\mathbf{g}},!}(n)$ of $\Lambda^{\tilde{\mathbf{g}}}(n)$ is the n-Auslander absolutely n-complete algebra $T_m^{(n)}(k)$.*

It is easy to see that the set of the projective vertices in $Q(n)$ is $\mathcal{P} = \{\mathbf{i} | i_n = 1\}$, and the set of the injective vertices is $\mathcal{I} = \{\mathbf{i} |\, |\mathbf{i}| = n+m-1\}$.

Let $m \ge 3$ and let $Q(n)$ be an n-cubic pyramid quiver. Define *stable n-cubic pyramid quiver* $\tilde{Q}(n)$ of height m as the quiver with

$$\begin{array}{ll} \text{vertex set:} & \tilde{Q}(n)_0 = Q(n)_0 \\ \text{arrow set:} & \tilde{Q}(n)_1 = Q(n)_1 \cup \{\gamma_{\mathbf{i}}^{(n+1)} : \mathbf{i} \longrightarrow \mathbf{i} - \mathbf{e}_n | i_n > 1\}. \end{array} \tag{6}$$

The stable quiver of our early example is as follow:



It follows from [14] and [12] that such quiver is a truncation of the McKay quiver of some finite Abelian subgroup of $SL_{n+1}(\mathbb{C})$.

Let $\bar{\mathbf{g}} = (\mathbf{g}, g')$ for some graded endomorphism g' induced by the $(n-1)$-translation defined in Theorem 2.1, then $g'(\gamma^{(s)}_{\mathbf{i}-\mathbf{e}_n}) = d_{n+1,s,\mathbf{i}-\mathbf{e}_n}\gamma^{(s)}_{\mathbf{i}-\mathbf{e}_n}$ for some $0 \neq d_{n+1,s,\mathbf{i}-\mathbf{e}_n} \in k$, since there is at most one arrow of each type s from any vertex of $Q(n)$. Define a relation set

$$\begin{array}{rcl} \tilde{\rho}^{\bar{\mathbf{g}}}(n) & = & \rho^{\mathbf{g}} \cup \{\gamma^{(n+1)}_{\mathbf{i}-\mathbf{e}_n}\gamma^{(n+1)}_{\mathbf{i}} \mid i_n > 1\} \\ & & \cup\{d_{n+1,s,\mathbf{i}-\mathbf{e}_n}\gamma^{(s)}_{\mathbf{i}-\mathbf{e}_n}\gamma^{(n+1)}_{\mathbf{i}} - \gamma^{(n+1)}_{\mathbf{i}-\mathbf{e}_{s-1}+\mathbf{e}_s}\gamma^{(s)}_{\mathbf{i}} \mid \mathbf{i} \in \tilde{Q}(n)_0, \\ & & \quad i_n > 1, i_{s-1} > 1 \text{ and } \sum_{t=1}^{s} i_t < m+s-1, 1 \leq s \leq n\}. \end{array} \tag{7}$$

Let $\tilde{\Lambda}^{\bar{\mathbf{g}}}(n)$ be the algebra given by the stable pyramid n-cubic quiver $\tilde{Q}(n)$ and the relation $\tilde{\rho}^{\bar{\mathbf{g}}}(n)$. Our key results in this paper, Lemma 3.3, essentially proves the following theorem.

Theorem 2.3. *$\tilde{\Lambda}^{\bar{\mathbf{g}}}(n)$ is an n-translation algebra with stable n-translation quiver and trivial n-translation.*

Set $\mathcal{R} = \{1, 2, \ldots, n\}$ and $\mathcal{R}^+ = \{1, 2, \ldots, n+1\}$. Write

$$\gamma^{(t_1,\ldots,t_s)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})} = \gamma^{(t_s)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}+\mathbf{e}_{t_1,\ldots,t_{s-1}})} \cdots \gamma^{(t_1)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})}$$

for a path determined by a sequences $\{t_1, \ldots, t_s\} \subset \mathcal{R}^+$. For a subset $P = \{p_1, \ldots, p_s\}$ of $\mathcal{R}^+$ with $1 \leq p_1 < \cdots < p_s \leq n+1$, write

$$\gamma^{(P)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})} = \gamma^{(p_1,\ldots,p_s)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})}.$$

We obviously have the following result.

Lemma 2.4. *Assume that $0 \leq l \leq n+1$. For any two vertices $\mathbf{i}, \mathbf{j} \in \tilde{Q}(n)_0$, we have that $\dim_k e_{\mathbf{j}}\tilde{\Lambda}^{\bar{\mathbf{g}}}(n)_l e_{\mathbf{i}} \leq 1$, and $\dim_k e_{\mathbf{j}}\tilde{\Lambda}^{\bar{\mathbf{g}}}(n)_l e_{\mathbf{i}} = 1$ if and only if any path of length l from $\mathbf{i}$ to $\mathbf{j}$ consists of arrows of different types.*

So we see that a bound path starting at a vertex $\mathbf{i}$ in $\tilde{Q}(n)$ is a path on $H^{\mathbf{i}}$ starting from $\mathbf{i}$. Since the n-translation is defined by $\tau\mathbf{i} = \mathbf{i} - \mathbf{e}_n$ for each non-projective vertex $\mathbf{i}$ in $Q(n)_0$, that is, the ones with $\mathbf{i}_n > 1$.

The following theorem follows from Theorem 5.2 and Proposition 4.2 of [15].

Theorem 2.5. *$\tilde{\Lambda}^{\bar{\mathbf{g}}}(n)$ is isomorphic to a twisted trivial extension of $\Lambda(n)$.*

For any $\mathbf{i} \in Q(n)_0$, let $S(\mathbf{i})$ be the simple $\Lambda(n)$-module at $\mathbf{i}$ and let $P(\mathbf{i})$ and $I(\mathbf{i})$ be its projective cover and injective envelop as $\Lambda(n)$-modules, respectively. Let $\tilde{S}(\mathbf{i})$ be the simple $\tilde{\Lambda}(n)$-module at $\mathbf{i}$ and let $\tilde{P}(\mathbf{i})$ and $\tilde{I}(\mathbf{i})$ be its projective cover and injective envelop as $\tilde{\Lambda}(n)$-modules, respectively. Conventionally, set $\tilde{S}(\mathbf{i})$, $\tilde{P}(\mathbf{i})$ and $\tilde{I}(\mathbf{i})$ to be zero when $\mathbf{i} \notin \tilde{Q}(n)_0$.

The supports of $\tilde{P}(\mathbf{i})$ and $\tilde{I}(\mathbf{i})$ are described with $U^{(n+1)}$, the following result follows directly from Lemma 2.4.

Lemma 2.6.

$$\mathbf{r}^t\tilde{P}(\mathbf{i})/\mathbf{r}^{t+1}\tilde{P}(\mathbf{i}) \simeq \bigoplus_{\mathbf{a}\in U_t^{(n+1)}} \tilde{S}(\mathbf{v}^{\mathbf{i}}(\mathbf{a})).$$

$$\operatorname{soc}^{t+1}\tilde{P}(\mathbf{i})/\operatorname{soc}^{t}\tilde{P}(\mathbf{i}) \simeq \bigoplus_{\mathbf{a}\in U_{n-t}^{(n+1)}} \tilde{S}(\mathbf{v}^{\mathbf{i}-\mathbf{e}_n}(\mathbf{e}-\mathbf{a}))$$

Since the components of $\mathbf{a}$ are 0 or 1, it is easy to see that $\mathbf{v}^{\mathbf{i}}(\mathbf{a})$ is a quiver vertex if and only if the following hold for all s: (i). $a_{s+1} \leq a_s$ if $i_s = 1$; (ii). $a_{s+1} \geq a_s$ if $\sum\limits_{t=1}^{s} i_t = m+s-1$. A vertex $\mathbf{i}$ with $i_s = 1$ or $\sum\limits_{t=1}^{s} i_t = m+s-1$ for some s is called a em boundary vertex, otherwise it is called an *internal vertex.* Now let $Z(\mathbf{i}) = \{s \in \mathcal{R} | i_s = 1\}$ and $W(\mathbf{i}) = \{s \in \mathcal{R} | \sum\limits_{t=1}^{s} i_t = m+s-1\}$.

The set $U^{(n+1)}(\mathbf{i})$ of $\mathbf{i}$-quiver vertices in $U^{(n+1)}$ is

$$U^{(n+1)}(\mathbf{i}) = \{\mathbf{a} = (a_1, \dots, a_{n+1}) | a_s \geq a_{s+1} \text{ if } s \in Z(\mathbf{i}), a_s \leq a_{s+1} \text{ if } s \in W(\mathbf{i})\}.$$

This implies that if $i_s = 1$ and $a_{s+1} = 1$, then $a_s = 1$, and if $s \in W(\mathbf{i})$ and $a_s = 1$, then $a_{s+1} = 1$. We can refine Lemma 2.6 as following

Lemma 2.7.

$$\mathbf{r}^t\tilde{P}(\mathbf{i})/\mathbf{r}^{t+1}\tilde{P}(\mathbf{i}) \simeq \bigoplus_{\mathbf{a}\in U_t^{(n+1)}(\mathbf{i})} \tilde{S}(\mathbf{v}^{\mathbf{i}}(\mathbf{a})).$$

$$\operatorname{soc}^{t+1}\tilde{P}(\mathbf{i})/soc^{t}\tilde{P}(\mathbf{i}) \simeq \bigoplus_{\mathbf{a}\in U_{n-t}^{(n+1)}(\mathbf{i}-\mathbf{e}_n)} \tilde{S}(\mathbf{v}^{\mathbf{i}-\mathbf{e}_n}(\mathbf{e}-\mathbf{a}))$$

Now consider $\overline{Q}(n) = \mathbb{Z}|_{(n-1)}Q(n)$ and 0-extension $\overline{\Lambda}(n)^{\bar{\mathbf{g}}} = \tilde{\Lambda}^{\bar{\mathbf{g}}}(n)\#\mathbb{Z}^*$, respectively (See [15]). We have

$$\begin{aligned}
&\overline{Q}(n)_0 = Q(n)_0 \times \mathbb{Z}\\
&\overline{Q}(n)_1 = Q(n)_1 \times \mathbb{Z} \cup \{\gamma_{\mathbf{i},v}^{(n+1)} : (\mathbf{i}, v) \longrightarrow (\mathbf{i}-\mathbf{e}_n, v+1) | (\mathbf{i}, v), (\mathbf{i}-\mathbf{e}_n, v+1) \in \overline{Q}(n)_0\}.
\end{aligned} \tag{8}$$

Set $\mathbf{e}_t^{(n+1)} = (\mathbf{e}_t^{(n)}, 0)$ for $1 \leq t \leq n$ and $\mathbf{e}_{n+1}^{(n+1)} = (\mathbf{0}^{(n)}, 1)$, then the arrows are $\gamma_{\mathbf{i}}^{(t)} : \mathbf{i} \longrightarrow \mathbf{i} - \mathbf{e}_{t-1} + \mathbf{e}_t$ for $\mathbf{i} \in \overline{Q}(n)_0$, $1 \leq t \leq n+1$. And the relations for $\overline{\Lambda}^{\bar{\mathbf{g}}}(n)$ is induced from $\tilde{\rho}^{\bar{\mathbf{g}}}(n)$. Thus

$$\begin{aligned}
\bar{\rho}^{\bar{\mathbf{g}}}(n) \quad = \quad & \rho^{\mathbf{g}} \times \mathbb{Z} \cup \{\gamma_{\mathbf{i}-\mathbf{e}_n+\mathbf{e}_{n+1}}^{(n+1)} \gamma_{\mathbf{i}}^{(n+1)} | i_n > 1\}\\
& \cup \{d_{n+1,s,\mathbf{i}-\mathbf{e}_n+\mathbf{e}_{n+1}} \gamma_{\mathbf{i}-\mathbf{e}_n}^{(s)} \gamma_{\mathbf{i}}^{(n+1)} - \gamma_{\mathbf{i}-\mathbf{e}_{s-1}+\mathbf{e}_s}^{(n+1)} \gamma_{\mathbf{i}}^{(s)} |\\
& \qquad \mathbf{i} \in \overline{Q}(n)_0, i_n > 1, i_{s-1} > 1 \text{ and } \sum_{t=1}^{s} i_t < m+s-1, 1 \leq s \leq n\}.
\end{aligned} \tag{9}$$

The following theorem follows from Theorem 2.1, Theorem 2.5 and Theorem 5.3 of [15].

Theorem 2.8. *$\overline{\Lambda}^{\bar{\mathbf{g}}}(n)$ is an n-translation algebra with stable n-translation quiver $\overline{Q}(n)$ and n-translation $\overline{\tau}_n : \mathbf{i} = \mathbf{i} - \mathbf{e}_{n+1}$.*

Let (Q,ρ) be a bound quiver and Q' a full subquiver of Q. Let $\rho[Q'] = \{e_j \sum a_p p e_i | \sum a_p p \in \rho, i, j \in Q'_0\}$, we call $(Q', \rho[Q'])$ a *full bound subquiver of* Q if for any $i, j \in Q'_0$, and for any path p in Q with $a_p \neq 0$ for some $\sum a_p p \in \rho$ and $e_j p e_i \neq 0$, then p is a path in Q'. In this case, we say $\rho' = \rho[Q']$ is a *restriction* of ρ on Q'.

Obviously, we have the following lemma.

Lemma 2.9. *Let $\Lambda = kQ/(\rho)$ be the algebra with bound quiver (Q,ρ) and let (Q',ρ') be a full bound subquiver of (Q,ρ). Let I be the ideal generated by the set of idempotents $\{e_j | j \in Q_0 \setminus Q'_0\}$. Then*

$$kQ'/(\rho') \simeq \Lambda/I.$$

3. Minimal Projective Resolutions of the Simples of $\tilde{\Lambda}^{\bar{\mathbf{g}}}(n)$

In this section, we study the projective resolution of the simples of the algebra $\tilde{\Lambda}^{\bar{\mathbf{g}}}(n)$. Fix $\mathbf{i} \in \tilde{Q}(n)_0$, the following lemma follows from the definition of $\mathbf{v}^{\mathbf{i}}(\mathbf{a})$ for $\mathbf{a} \in \mathbb{Z}^{+}{}^{n+1}$.

Lemma 3.1. *For $0 \leq l \leq m-1$, if $\mathbf{a}, \mathbf{a}' \in \mathbb{Z}^{+}{}^{n+1}_{l}$, then $\mathbf{v}^{\mathbf{i}}(\mathbf{a}) = \mathbf{v}^{\mathbf{i}}(\mathbf{a}')$ if and only if $\mathbf{a} = \mathbf{a}'$.*

Fix $\mathbf{a} \in \mathbb{Z}^{+}{}^{n+1}$. Let $O(\mathbf{a}) = \{t | a_t = 0\}$, $T(\mathbf{a}) = \{1 \leq t \leq n+1 | a_t > b_t\}$ and let $\mathcal{R}(\mathbf{a}) = \mathcal{R}^+ \setminus (O(\mathbf{a}) \cup T(\mathbf{a}))$ for any $\mathbf{a} \in \mathbb{Z}^{+}{}^{n+1}$. Then $\mathbf{a} \in C^{(n)}(\mathbf{i})$ if and only if $T(\mathbf{a}) = \emptyset$.

Let

$$\cdots \longrightarrow \tilde{P}^2(\tilde{S}(\mathbf{i})) \xrightarrow{f_2} \tilde{P}^1(\tilde{S}(\mathbf{i})) \xrightarrow{f_1} \tilde{P}^0(\tilde{S}(\mathbf{i})) \xrightarrow{f_0} \tilde{S}(\mathbf{i}) \longrightarrow 0 \tag{10}$$

be a minimal projective resolution of the simple $\tilde{\Lambda}$-modules $\tilde{S}(\mathbf{i})$ corresponding to the vertex $\mathbf{i} \in \tilde{Q}(n)_0$.

Our main aim of this section is characterizing the linear part of this projective resolution using $(n+1)$-cuboid $C^{(n)}(\mathbf{i})$ as in the following proposition. The technical detail of the proof will be given in Lemma 3.3. For the last assertion, notice the fact that $\mathbf{v}^{\mathbf{i}}(\mathbf{a} + \mathbf{e}) = \mathbf{v}^{\mathbf{i}}(\mathbf{a})$.

Proposition 3.2. *For $l = 0, 1, \ldots, m-1$*

$$\tilde{P}^l(\tilde{S}(\mathbf{i})) \simeq \bigoplus_{\mathbf{a} \in C^{(n)}_l(\mathbf{i})} \tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a})), \tag{11}$$

and $\operatorname{Ker} f_{m-1} \simeq \tilde{S}(\mathbf{v}^{\mathbf{i}}(\mathbf{b}(\mathbf{i})))$.

Let $\{\epsilon^{(l)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})} | \mathbf{a} \in C^{(n)}_l(\mathbf{i})\}$ be the standard basis of the direct sum (11), then

$$e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})} \epsilon^{(l)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})} = \epsilon^{(l)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})} \text{ and } e_{\mathbf{i}'} \epsilon^{(l)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})} = 0 \text{ if } \mathbf{i}' \neq \mathbf{v}^{\mathbf{i}}(\mathbf{a}).$$

We assume that $\epsilon^{(l)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})} = 0$ when $\mathbf{a} \notin C^{(n)}_l(\mathbf{i})$ or $\mathbf{v}^{\mathbf{i}}(\mathbf{a}) \notin \tilde{Q}(n)_0$, for $l = 1, \ldots, m-1$, conventionally.

Lemma 3.3. *In the projective resolution* (10)*, we have*

(a) (11) *hold for* $0 \le l \le m-1$*;*

(b) *for* $1 \le l \le m-1$*, and for each* $\mathbf{a} \in C_l^{(n)}(\mathbf{i})$*, there are elements* $\zeta^{(l-1)}_{\mathbf{a}-\mathbf{e}_t} \in k$ *for* $t = 1, \dots, n+1$*, with* $\zeta^{(l-1)}_{\mathbf{a}-\mathbf{e}_t} \neq 0$ *if* $\mathbf{a}-\mathbf{e}_t \in C_{l-1}^{(n)}(\mathbf{i})$*, and* $\zeta^{(l-1)}_{\mathbf{a}-\mathbf{e}_t} = 0$ *otherwise, such that the element* $\theta^{(l-1)}_{\mathbf{v^i}(\mathbf{a})} = \sum\limits_{t=1}^{n+1} \zeta^{(l-1)}_{\mathbf{v^i}(\mathbf{a}-\mathbf{e}_t)} \gamma^{(t)}_{\mathbf{v^i}(\mathbf{a}-\mathbf{e}_t)} \epsilon^{(l)}_{\mathbf{v^i}(\mathbf{a}-\mathbf{e}_t)}$ *satisfies*

$$e_{\mathbf{v^i}(\mathbf{a})} \theta^{(l-1)}_{\mathbf{v^i}(\mathbf{a})} = \theta^{(l-1)}_{\mathbf{v^i}(\mathbf{a})} \tag{12}$$

and the map f_l *is defined by*

$$f_l(\epsilon^{(l)}_{\mathbf{v^i}(\mathbf{a})}) = \theta^{(l-1)}_{\mathbf{v^i}(\mathbf{a})}; \tag{13}$$

(c)

$$K_l = \{\theta^{(l)}_{\mathbf{v^i}(\mathbf{a})} | \mathbf{a} \in C_{l+1}^{(n)}(\mathbf{i})\} \tag{14}$$

generates Ker f_l *for* $0 \le l \le m-2$*;*

(d) $C_{m-1}^{(n)}(\mathbf{i}) = \{\mathbf{b}\}$*,* $\mathbf{b}+\mathbf{e}$ *is a* $\mathbf{i}$*-quiver vertex and* Ker $f_{m-1} \simeq S(\mathbf{v^i}(\mathbf{b}+\mathbf{e}))$*.*

Proof. We now prove (a), (b) and (c) using induction on l. The assertions for $\tilde{P}^0(S(\mathbf{i})), \operatorname{Ker} f_0$ and f_1 are obvious.

Assume that $1 \le l \le m-1$ and the assertions hold for $\tilde{P}^h(\mathbf{i}), \operatorname{Ker} f_h$, and f_h for $h < l$. By the inductive assumption, Ker f_{l-1} is a graded module generated in degree l and as k-spaces

$$(\operatorname{Ker} f_{l-1})_l = \operatorname{top}(\operatorname{Ker} f_{l-1}) = L(\theta^{(l-1)}_{\mathbf{v^i}(\mathbf{a})} | \mathbf{a} \in C_l^{(n)}(\mathbf{i})) = \sum_{\mathbf{a} \in C_l^{(n)}(\mathbf{i})} k\theta^{(l-1)}_{\mathbf{v^i}(\mathbf{a})}.$$

By (12), we have

$$k\theta^{(l-1)}_{\mathbf{v^i}(\mathbf{a})} \simeq \tilde{\Lambda}_0 e_{\mathbf{v^i}(\mathbf{a})}, \tag{15}$$

as $\tilde{\Lambda}_0$-modules. Since for each $\mathbf{a} \in C_l^{(n)}(\mathbf{i})$, we have exactly one nonzero element $\theta^{(l-1)}_{\mathbf{v^i}(\mathbf{a})}$ in K_{l-1}. Thus

$$\operatorname{top}(\operatorname{Ker} f_{l-1}) \simeq \bigoplus_{\mathbf{a} \in C_l^{(n)}(\mathbf{i})} \tilde{\Lambda}_0 e_{\mathbf{v^i}(\mathbf{a})}.$$

Thus

$$\tilde{P}^l(S(\mathbf{i})) \simeq \bigoplus_{\mathbf{a} \in C_l^{(n)}(\mathbf{i})} \tilde{P}(\mathbf{v^i}(\mathbf{a})).$$

This proves (a) for l.

Define the homomorphism f_l from $\tilde{P}^l(S(\mathbf{i}))$ to $\tilde{P}^{l-1}(S(\mathbf{i}))$ with

$$f_l(\epsilon^{(l)}_{\mathbf{v^i}(\mathbf{a})}) = \theta^{(l-1)}_{\mathbf{v^i}(\mathbf{a})},$$

Clearly f_l is an epimorphism from $\tilde{P}^l(S(\mathbf{i}))$ to Ker f_{l-1}, and $f_l(\tilde{P}^l(S(\mathbf{i})))_p \subseteq (\tilde{P}^{l-1}(S(\mathbf{i})))_{p+1}$ for $p \in \mathbb{Z}$. This proves (13) for l.

Now we compute generators of Ker f_l for $l \le m-2$. Clearly $(\operatorname{Ker} f_l)_l = 0$ and $(\operatorname{Ker} f_l)_{l+n+1} = \operatorname{soc} \tilde{P}^l(S(\mathbf{i}))$, by the inductive assumption.

Assume that $0 \neq x_p$ is a homogeneous element in $(\operatorname{Ker} f_l)_p$, then we can write $x_{l+1} = \sum\limits_{\mathbf{a} \in C_l^{(n)}(\mathbf{i})} (\sum\limits_{t=1}^{n+1} z_{\mathbf{a},t} \gamma^{(t)}_{\mathbf{v^i}(\mathbf{a})}) \epsilon^{(l)}_{\mathbf{v^i}(\mathbf{a})}$, with the convention that $z_{\mathbf{a},t} = 0$ provided that $\mathbf{v}^{\mathbf{i}}(\mathbf{a}) \notin \tilde{Q}(n)_0$. Thus

$$\begin{array}{rcl} 0 & = & f_l(x_{l+1}) = \sum\limits_{\mathbf{a}\in C_l^{(n)}(\mathbf{i})} (\sum\limits_{t=1}^{n+1} z_{\mathbf{a},t}\gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})}) f_l(\epsilon^{(l)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})}) \\ & = & \sum\limits_{\mathbf{a}\in C_l^{(n)}(\mathbf{i})} (\sum\limits_{t=1}^{n+1} z_{\mathbf{a},t}\gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})}) \cdot \theta^{(l-1)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})} \\ & = & \sum\limits_{\mathbf{a}\in C_l^{(n)}(\mathbf{i})} (\sum\limits_{t=1}^{n+1} z_{\mathbf{a},t}\gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})}) \cdot (\sum\limits_{t=1}^{n+1} \zeta^{(l-1)}_{\mathbf{a}-\mathbf{e}_t}\gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}-\mathbf{e}_t)}\epsilon^{(l-1)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}-\mathbf{e}_t)}) \\ & = & \sum\limits_{\mathbf{a}\in C_l^{(n)}(\mathbf{i})} (\sum\limits_{\substack{t'\neq t \\ 1\le t,t'\le n+1}} z_{\mathbf{a},t}\zeta^{(l-1)}_{\mathbf{a}-\mathbf{e}_{t'}} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}+\mathbf{e}_t)}\gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})}\gamma^{(t')}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}-\mathbf{e}_{t'})}\epsilon^{(l-1)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}-\mathbf{e}_{t'})}) \end{array} \tag{16}$$

Assume first that $\mathbf{a}' \in {\mathbb{Z}^+}^{n+1}_{l+1}$ and $\mathbf{a}'-\mathbf{e}_t, \mathbf{a}'-\mathbf{e}_{t'} \in C_l^{(n)}(\mathbf{i})$ for $t \neq t'$. Then $0 \le a'_t = (\mathbf{a}'-\mathbf{e}_{t'})_t \le b_t$, $0 \le a'_{t'} = (\mathbf{a}'-\mathbf{e}_t)_{t'} \le b_{t'}$, and $0 \le a_{t''} = (\mathbf{a}'-\mathbf{e}_t)_{t''} \le b_{t''}$ for $t'' \notin \{t,t'\}$. Thus $\mathbf{a}' \in C_{l+1}^{(n)}(\mathbf{i})$, and we have that if $\mathbf{a}'-\mathbf{e}_t-\mathbf{e}_{t'} \in C^{(n)}(\mathbf{i})$, then $\mathbf{a}'-\mathbf{e}_t, \mathbf{a}'-\mathbf{e}_{t'} \in C^{(n)}(\mathbf{i})$, left multiple (16) with $e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}$, we get,

$$\begin{array}{rcl} 0 & = & \sum_{1\le t<t'\le n+1, \mathbf{a}'-\mathbf{e}_t-\mathbf{e}_{t'}\in C^{(n)}(\mathbf{i})} \\ & & (z_{\mathbf{a}'-\mathbf{e}_t,t}\zeta^{(l-1)}_{\mathbf{a}'-\mathbf{e}_t-\mathbf{e}_{t'},t'} - z_{\mathbf{a}'-\mathbf{e}_{t'},t'}\zeta^{(l-1)}_{\mathbf{a}'-\mathbf{e}_{t'}-\mathbf{e}_t,t} d_{t',t,\mathbf{a}'-\mathbf{e}_{t'}-\mathbf{e}_t}) \\ & & e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}\gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_t)}\gamma^{(t')}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_t-\mathbf{e}_{t'})}\epsilon^{(l-1)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_t-\mathbf{e}_{t'})}. \end{array} \tag{17}$$

So we get a system of linear equations

$$z_{\mathbf{a}'-\mathbf{e}_t,t}\zeta^{(l-1)}_{\mathbf{a}'-\mathbf{e}_t-\mathbf{e}_{t'},t'} - z_{\mathbf{a}'-\mathbf{e}_{t'},t'}\zeta^{(l-1)}_{\mathbf{a}'-\mathbf{e}_{t'}-\mathbf{e}_t,t} d_{t',t,\mathbf{a}'-\mathbf{e}_{t'}-\mathbf{e}_t} \quad = \quad 0 \tag{18}$$

for $1 \le t < t' \le n+1$ with $\mathbf{a}'-\mathbf{e}_t-\mathbf{e}_{t'} \in C_{l-1}^{(n)}(\mathbf{i})$.

For a path p ending at $\mathbf{v}^{\mathbf{i}}(\mathbf{a}')$, the types of the arrows in p are in

$$\{1,\ldots,n,n+1\} \setminus O(\mathbf{a}').$$

Let $l' = l-n-1+|O(\mathbf{a}')|$, the composition factor $\tilde{S}(\mathbf{a}')$ in the degree $l+1$ components of the minimal projective resolution (10) is

$$\begin{array}{rcl} 0 & \longrightarrow & e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\operatorname{Ker} f_l)_{l+1} \longrightarrow \bigoplus_{\mathbf{a}\in C_l^{(n)}(\mathbf{i})} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}))[l])_{l+1} \\ & & \longrightarrow \bigoplus_{\mathbf{a}\in C_{l-1}^{(n)}(\mathbf{i})} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}))[l-1])_{l+1} \longrightarrow \cdots \\ & & \longrightarrow \bigoplus_{\mathbf{a}\in C_{l'}^{(n)}(\mathbf{i})} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}))[l'])_{l+1} \longrightarrow 0. \end{array} \tag{19}$$

Note that

$$\bigoplus_{\mathbf{a}\in C_{t'}^{(n)}(\mathbf{i})} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}))[t'])_{l+1} = \bigoplus_{\mathbf{a}\in C_{t'}^{(n)}(\mathbf{i})} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}\mathbf{r}^{l+1-t'}\tilde{\Lambda}(n)/\mathbf{r}^{l+2-t'}\tilde{\Lambda}(n)e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})}$$

is the space spanned by the bound paths of length $l+1-t'$ ending at $\mathbf{v}^{\mathbf{i}}(\mathbf{a}')$. Since bound paths are formed by arrows of different types, and two bound paths are linearly independent if and only if the set of the types of their arrows are different, a maximal linearly independent set of paths of length t ending at $\mathbf{v}^{\mathbf{i}}(\mathbf{a}')$ has $\begin{pmatrix} n+1-|O(\mathbf{a}')| \\ t \end{pmatrix}$ elements, that is

$$\dim_k \bigoplus_{\mathbf{a}\in C_{t'}^{(n)}(\mathbf{i})} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}))[t'])_{l+1} = \begin{pmatrix} n+1-|O(\mathbf{a}')| \\ l+1-t' \end{pmatrix}.$$

So it follows from (19) that

$$\begin{array}{rcl} 0 &=& \dim{}_k e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\operatorname{Ker} f_l)_{l+1} \\ && + \sum\limits_{t=1}^{n+1-|O(\mathbf{a}')|} (-1)^t \dim{}_k \bigoplus_{\mathbf{a}\in C^{(n)}_{l-t+1}(\mathbf{i})} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}))[l-t+1])_{l+1} \\ &=& \dim{}_k e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\operatorname{Ker} f_l)_{l+1} + \sum\limits_{t=1}^{n+1-|O(\mathbf{a}')|} (-1)^t \left(\begin{array}{c} n+1-|O(\mathbf{a}')| \\ t \end{array} \right) \\ &=& \dim{}_k e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\operatorname{Ker} f_l)_{l+1} - 1 + (1-1)^{n+1-|O(\mathbf{a}')|}. \end{array}$$

Thus $\dim{}_k e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\operatorname{Ker} f_l)_{l+1} = 1$, and this is equivalent to that the solution space of (18) is one dimensional. Take a nonzero solution

$$z_{\mathbf{a}'-\mathbf{e}_t,t} = \zeta^{(l)}_{\mathbf{a}'-\mathbf{e}_t,t} \in k,$$

for $\mathbf{a}' \in C^{(n)}_{l+1}(\mathbf{i})$ with $\mathbf{a}' - \mathbf{e}_{t''} - \mathbf{e}_{t'} \in C^{(n)}(\mathbf{i})$ for some $1 \le t' < t'' \le n+1$. Then each $\zeta^{(l)}_{\mathbf{a}'-\mathbf{e}_t,t} \ne 0$ if $a'_t \ne 0$, and we take $\zeta^{(l)}_{\mathbf{a}'-\mathbf{e}_t,t} = 0$ for those t with $a'_t = 0$.

$$\theta^{(l)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')} = \sum_{t=1}^{n+1} \zeta^{(l)}_{\mathbf{a}'-\mathbf{e}_t,t} \gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_t)} \epsilon^{(l)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_t)} \tag{20}$$

is a nonzero element of $e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\operatorname{Ker} f_l)_{l+1}$.

Assume now that $\mathbf{a}' \in {\mathbb{Z}^+}^{n+1}_{l+1}$, $\mathbf{a}' - \mathbf{e}_t \in C^{(n)}_l(\mathbf{i})$ and $\mathbf{a}' - \mathbf{e}_{t'} \notin C^{(n)}_l(\mathbf{i})$ for $t' \ne t$.

If $\mathbf{a}' \in C^{(n)}_{l+1}(\mathbf{i})$, then $a'_{t'} = 0$ for $t' \ne t$. The exact sequence (19) of composition factor $\tilde{S}(\mathbf{a}')$ in the degree $l+1$ component of the projective resolution (10) becomes

$$0 \longrightarrow e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\operatorname{Ker} f_l)_{l+1} \longrightarrow e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_t))[l])_{l+1} \longrightarrow 0$$

and $\dim{}_k e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\operatorname{Ker} f_l)_{l+1} = 1$. Left multiple (16) with $e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}$, we get a zero term:

$$z_{\mathbf{a}'-\mathbf{e}_t,t} \zeta^{(l-1)}_{\mathbf{a}'-\mathbf{e}_t-\mathbf{e}_t,t} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')} \gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_t)} \gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_t-\mathbf{e}_t)} \epsilon^{(l-1)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_t-\mathbf{e}_t)} = 0.$$

So we can take $z_{\mathbf{a}'-\mathbf{e}_t,t} = 1$ and

$$\theta^{(l)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')} = \gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_t)} \epsilon^{(l)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_t)}$$

is a nonzero element of $e_{\mathbf{a}'}(\operatorname{Ker} f_l)_{l+1}$.

Now we consider the case that $\mathbf{a}' \in {\mathbb{Z}^+}^{n+1}_{l+1} \setminus C^{(n)}_{l+1}(\mathbf{i})$ such that $\mathbf{a}' - \mathbf{e}_t \in C^{(n)}_l(\mathbf{i})$ and $\mathbf{a}' - \mathbf{e}_{t'} \notin C^{(n)}_l(\mathbf{i})$ for $t' \ne t$. In this case $a'_t = b_t(\mathbf{i}) + 1$. We show that $\tilde{S}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}'))$ is not a composition factor in $(\operatorname{Ker} f_l)_{l+1}$

Assume that $\mathbf{v}^{\mathbf{i}}(\mathbf{a}') \in \tilde{Q}(n)_0$ and consider the composition factor $\tilde{S}(\mathbf{a}')$ in the degree $l+1$ component of the projective resolution (10). Since in this case, each bound path from a vertex $\mathbf{v}^{\mathbf{i}}(\mathbf{a})$ with $\mathbf{a} \in C^{(n)}(\mathbf{i})$ to $\mathbf{v}^{\mathbf{i}}(\mathbf{a}')$ in $\tilde{Q}(n)$ is a multiple of a path of same length with the last arrow $\gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_t)}$, so (19) becomes

$$\begin{array}{rcl} 0 &\longrightarrow& e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\operatorname{Ker} f_l)_{l+1} \longrightarrow \bigoplus_{\mathbf{a}\in C^{(n)}_l(\mathbf{i})} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')} \gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_t)} (\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}))[l])_l \longrightarrow \cdots \\ &\longrightarrow& \bigoplus_{\mathbf{a}\in C^{(n)}_{l'}(\mathbf{i})} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')} \gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_t)} (\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}))[l'])_l \longrightarrow 0. \end{array}$$

Similar to the argument after (19), we get that $e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\operatorname{Ker} f_l)_{l+1} = 0$.

This proves that the set K_l spans the space $(\operatorname{Ker} f_l)_{l+1}$.

Since f_l is degree zero map, $\operatorname{Ker} f_l$ is graded. Consider the degree $l+s$ component of (10), for each $2 \le s \le n+1$, we have for each $\mathbf{a}' \in {\mathbb{Z}^+}^{n+1}_{l+s}$,

$$\begin{array}{rcl} 0 &\longrightarrow& e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\operatorname{Ker} f_l)_{l+s} \longrightarrow \bigoplus_{\mathbf{a}\in C^{(n)}_l(\mathbf{i})} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}))[l])_{l+s} \longrightarrow \cdots \\ &\longrightarrow& \bigoplus_{\mathbf{a}\in C^{(n)}_{l'}(\mathbf{i})} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}))[l''])_{l+s} \longrightarrow 0, \end{array} \tag{21}$$

with $l'' = l+s-n-1+|O(\mathbf{a}')|$.

Note that $\bigoplus_{\mathbf{a}\in C^{(n)}_{l-s'}(\mathbf{i})} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}))[l-s'])_{l+s}$ is in the space of the bound paths of length $s+s'$ to the vertex $e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}$ in $\tilde{Q}(n)_0$. Thus paths formed by arrows of different types are nonzero and the representatives of those with the same set of types form a basis. On the other hand, since there is $\mathbf{a}'' \in C^{(n)}_l(\mathbf{i})$ and a bound path of length s from $\mathbf{v}^{\mathbf{i}}(\mathbf{a}'')$ to $\mathbf{v}^{\mathbf{i}}(\mathbf{a}')$, and $a''_{t'} \le b_{t'}$ for all t', thus $|T(\mathbf{a}')| \le s$. So for any $\mathbf{a} \in C^{(n)}(\mathbf{i})$, each bound path from $\mathbf{v}^{\mathbf{i}}(\mathbf{a})$ to $\mathbf{v}^{\mathbf{i}}(\mathbf{a}')$ passes the arrows of all the types in $T(\mathbf{a}')$. This implies that a set of the linear independent path to $\mathbf{v}^{\mathbf{i}}(\mathbf{a}')$ is determined by the arrows of types in $\mathcal{R}^+ \setminus (O(\mathbf{a}') \cup T(\mathbf{a}'))$. Thus

$$\dim{}_k \bigoplus_{\mathbf{a}\in C^{(n)}_{l-s'}(\mathbf{i})} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}))[l-s'])_{l+s} = \left(\begin{array}{c} n+1-|O(\mathbf{a}')|-|T(\mathbf{a}')| \\ s+s'-|T(\mathbf{a}')| \end{array}\right)$$

and

$$\begin{aligned}
& \dim{}_k e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\operatorname{Ker} f_l)_{l+s} \\
= & \sum_{s'=0}^{n+1-|O(\mathbf{a}')|-|T(\mathbf{a}')|-s} (-1)^{s'} \sum_{\mathbf{a}\in C^{(n)}_{l+s}(\mathbf{i})} \dim{}_k e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}))[l])_{l+s} \\
= & \sum_{s'=0}^{n+1-|O(\mathbf{a}')|-|T(\mathbf{a}')|-s} (-1)^{s'} \left(\begin{array}{c} n+1-|O(\mathbf{a}')|-|T(\mathbf{a}')| \\ s-|T(\mathbf{a}')|+s' \end{array}\right) \\
= & \left(\begin{array}{c} n-|O(\mathbf{a}')|-|T(\mathbf{a}')| \\ s-|T(\mathbf{a}')|-1 \end{array}\right).
\end{aligned}$$

Now we consider the submodule generated by K_l. For each subset P of $s-|T(\mathbf{a}')|-1$ elements in $\mathcal{R}^+ \setminus (O(\mathbf{a}') \cup T(\mathbf{a}'))$, let

$$\begin{aligned}
\kappa(P,\mathbf{a}) &= \gamma^{(T(\mathbf{a}'))}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_{T(\mathbf{a}')})}\gamma^{(P)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})}\theta^{(l)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})} = \gamma^{(T(\mathbf{a}'))}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_{T(\mathbf{a}')})}\gamma^{(P)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})}\sum_{t=1}^{n+1}\zeta^{(l)}_{\mathbf{a}-\mathbf{e}_t,t}\gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}-\mathbf{e}_t)}\epsilon^{(l)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}-\mathbf{e}_t)} \\
&= \sum_{t\in\mathcal{R}^+\setminus(O(\mathbf{a}')\cup T(\mathbf{a}')\cup P)} \zeta^{(l)}_{\mathbf{a}-\mathbf{e}_t,t}\gamma^{(T(\mathbf{a}'))}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}'-\mathbf{e}_{O(T)})}\gamma^{(P)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a})}\gamma^{(t)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}-\mathbf{e}_t)}\epsilon^{(l)}_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}-\mathbf{e}_t)},
\end{aligned}$$

for any $\mathbf{a} = \mathbf{a}' - \mathbf{e}_{T(\mathbf{a}')\cup P} \in C^{(n)}_{l+s-|P|-|T(\mathbf{a}')|}(\mathbf{i})$.

Let $s'' = l+s-|P|-|T(\mathbf{a}')|$. Fix a $t_0 \in \mathcal{R}(\mathbf{a}')$, and let $\mathcal{P}(s'',t_0) = \{P \subset \mathcal{R}^+ \setminus (O(\mathbf{a}') \cup T(\mathbf{a}') \cup \{t_0\}) | |P| = s''\}$. Assume that

$$\sum_{P\in\mathcal{P}(s'',t_0),\mathbf{a}\in C^{(n)}_{s''}(\mathbf{i})} d_{P,\mathbf{a}}\kappa(P,\mathbf{a}) = 0$$

for $d_{P,\mathbf{a}} \in k$, that is

$$\begin{aligned}
0 &= \sum_{\substack{P\in\mathcal{P}(s'',t_0)\\ \mathbf{a}\in C^{(n)}_{s''}(\mathbf{i})}} d_{P,\mathbf{a}}\kappa(P,\mathbf{a})\\
&= \sum_{\substack{P\in\mathcal{P}(s'',t_0)\\ \mathbf{a}\in C^{(n)}_{s''}(\mathbf{i})}} \sum_{t\in\mathcal{R}(\mathbf{a}')\setminus P} d_{P,\mathbf{a}}\zeta^{(l)}_{\mathbf{a}-\mathbf{e}_t,t}\gamma^{(T(\mathbf{a}'))}_{\mathbf{v^i}(\mathbf{a}'-\mathbf{e}_{T(\mathbf{a}')})}\gamma^{(P)}_{\mathbf{v^i}(\mathbf{a})}\gamma^{(t)}_{\mathbf{v^i}(\mathbf{a}-\mathbf{e}_t)}\epsilon^{(l)}_{\mathbf{v^i}(\mathbf{a}-\mathbf{e}_t)}\\
&= \sum_{\substack{\mathbf{a}\in C^{(n)}_{s''-1}(\mathbf{i})\\ P\in\mathcal{P}(s'',t_0)}} \sum_{\substack{t\in\mathcal{R}(\mathbf{a}')\\ t\notin P}} d_{P,\mathbf{a}+\mathbf{e}_t}\zeta^{(l)}_{\mathbf{a},t}\gamma^{(T(\mathbf{a}'))}_{\mathbf{v^i}(\mathbf{a}'-\mathbf{e}_{T(\mathbf{a})})}\gamma^{(P)}_{\mathbf{v^i}(\mathbf{a}+\mathbf{e}_t)}\gamma^{(t)}_{\mathbf{v^i}(\mathbf{a})}\epsilon^{(l)}_{\mathbf{v^i}(\mathbf{a})}\\
&= \sum_{\substack{\mathbf{a}\in C^{(n)}_{s''-1}(\mathbf{i})\\ t,t'\in\mathcal{R}(\mathbf{a}')\setminus\{t_0\}}} \sum_{\substack{P\subset\mathcal{R}(\mathbf{a}')\setminus\{t_0\}\\ |P|=s''-1}} \\
&\qquad (d_{P\cup\{t'\},\mathbf{a}+\mathbf{e}_t}h_{t',P}\zeta^{(l)}_{\mathbf{a},t} + d_{P\cup\{t\},\mathbf{a}+\mathbf{e}_{t'}}h_{t,P}c(t,t')\zeta^{(l)}_{\mathbf{a},t'})\\
&\qquad \gamma^{(T(\mathbf{a}'))}_{\mathbf{v^i}(\mathbf{a}'-\mathbf{e}_{T(\mathbf{a})})}\gamma^{(P)}_{\mathbf{v^i}(\mathbf{a}+\mathbf{e}_{t,t'})}\gamma^{(t')}_{\mathbf{v^i}(\mathbf{a}+\mathbf{e}_t)}\gamma^{(t)}_{\mathbf{v^i}(\mathbf{a})}\epsilon^{(l)}_{\mathbf{v^i}(\mathbf{a})}\\
&\quad + \sum_{\substack{\mathbf{a}\in C^{(n)}_{s''-1}(\mathbf{i})\\ P\subset\mathcal{P}(s'',t_0)}} d_{P,\mathbf{a}+\mathbf{e}_{t_0}}\zeta^{(l)}_{\mathbf{a},t_0}\gamma^{(T(\mathbf{a}'))}_{\mathbf{v^i}(\mathbf{a}'-\mathbf{e}_{T(\mathbf{a})})}\gamma^{(P)}_{\mathbf{v^i}(\mathbf{a}+\mathbf{e}_{t_0})}\gamma^{(t_0)}_{\mathbf{v^i}(\mathbf{a})}\epsilon^{(l)}_{\mathbf{v^i}(\mathbf{a})},
\end{aligned} \tag{22}$$

for some $h_{t',P}, h_{t,P}, c(t,t') \in k$. Thus $d_{P,\mathbf{a}} = 0$ for all $P \in \mathcal{P}(s'', v_0)$ and $\mathbf{a} \in C^{(n)}_{s''}(\mathbf{i})$ with $\mathbf{a} - \mathbf{e}_{t_0} \in C^{(n)}_{s''-1}(\mathbf{i})$. Since $a'_{t_0} > 0$, and if $e_{\mathbf{v^i}(\mathbf{a}')}\gamma^{(T(\mathbf{a}'))}_{\mathbf{v^i}(\mathbf{a}'-\mathbf{e}_{T(\mathbf{a})})}\gamma^{(P)}_{\mathbf{v^i}(\mathbf{a})}\epsilon^{(l)}_{\mathbf{v^i}(\mathbf{a})} \neq 0$, then $\mathbf{a} = \mathbf{a}' - \mathbf{e}_{T(\mathbf{a})\cup P}$ and $a_{t_0} = a'_{t_0} > 0$, thus $\mathbf{a} - \mathbf{e}_{t_0} \in C^{(n)}_{s''-1}(\mathbf{i})$ for all $\mathbf{a} \in C^{(n)}_{s''}(\mathbf{i})$.

This shows that $\{\kappa(P,\mathbf{a}) | P \in \mathcal{P}(s'', t_0), \mathbf{a} \in C^{(n)}_{s''}(\mathbf{i})\}$ is a linearly independent set and thus

$$\dim_k(\tilde{\Lambda}(n)K_l)_{l+s'} \geq \begin{pmatrix} n - |O(\mathbf{a}')| - |T(\mathbf{a}')| \\ s - |T(\mathbf{a}')| - 1 \end{pmatrix}.$$

Since $(\tilde{\Lambda}(n)K_l)_{l+s'} \subset (\operatorname{Ker} f_l)_{l+s'}$, this implies that $(\tilde{\Lambda}(n)K_l)_{l+s'} = (\operatorname{Ker} f_l)_{l+s'}$ for $s' = 2, \ldots, n+1$.

This proves that $\operatorname{Ker} f_l$ is generated by K_l for $0 \leq l \leq m-2$, and so we prove (a), (b) and (c), by induction.

Now we prove (d), and consider the case of $l = m-1$, note that $C^{(n)}_{m-1}(\mathbf{i}) = \{\mathbf{b} = (b_1, \ldots, b_{n+1})\}$, with $b_1 = m + n - 1 - \sum\limits_{t=1}^{n} i_t$, and $b_t = i_{t-1} - 1$ for $2 \leq t \leq n+1$.

Clearly, we have that $\operatorname{soc} \tilde{P}(\mathbf{v^i}(\mathbf{b})) = \tilde{P}(\mathbf{v^i}(\mathbf{b}))_{n+1} = (\operatorname{Ker} f_{m-1})_{n+1} \subset \operatorname{Ker} f_{m-1}$. For each $\mathbf{a}' \neq \mathbf{b} + \mathbf{e}$ such that $\tilde{S}(\mathbf{v^i}(\mathbf{a}'))$ is a composition factor of $\mathbf{r}\tilde{P}(\mathbf{v^i}(\mathbf{b}))_s$, for $1 \leq s \leq n$, $e_{\mathbf{v^i}(\mathbf{a}')}\tilde{\Lambda}e_{\mathbf{v^i}(\mathbf{b})} \neq 0$ and $\mathbf{a}' \in {\mathbb{Z}^+}^{n+1}_{m-1+s} \setminus C^{(n)}(\mathbf{i})$. Thus $|T(\mathbf{a}')| = s$ and there is a bound path $\gamma^{(T(\mathbf{a}')^\sigma)}_{\mathbf{b}}$ of the arrows of the types in $T(\mathbf{a}')$ from $\mathbf{v^i}(\mathbf{b})$ to $\mathbf{v^i}(\mathbf{a}')$. So paths of the form $\gamma^{(T(\mathbf{a})^\sigma)}_{\mathbf{b}}\gamma^{(P)}_{\mathbf{v^i}(\mathbf{b}-\mathbf{e}_P)}$ for $P \subset \mathcal{R}(\mathbf{a}')$ with $|P| = s'$ form a basis of $\bigoplus_{\mathbf{a}\in C^{(n)}_{m-1-s'}(\mathbf{i})} e_{\mathbf{v^i}(\mathbf{a}')}\tilde{P}(\mathbf{v^i}(\mathbf{a}))_{s+s'}$ for $0 \leq s' \leq n + 1 - |O(\mathbf{a}')| - |T(\mathbf{a}')|$.

Consider the composition factor $\tilde{S}(\mathbf{a}')$ in the degree $m+s$ component of the projective resolution (10), we get

$$\begin{aligned}
0 &\longrightarrow e_{\mathbf{v^i}(\mathbf{a}')}(\operatorname{Ker} f_{m-1})_{m-1+s} \longrightarrow e_{\mathbf{v^i}(\mathbf{a}')}(\tilde{P}(\mathbf{v^i}(\mathbf{b}))[m-1])_{m-1+s}\\
&\longrightarrow \textstyle\bigoplus_{\mathbf{a}\in C^{(n)}_{m-2}(\mathbf{i})} e_{\mathbf{v^i}(\mathbf{a}')}(\tilde{P}(\mathbf{v^i}(\mathbf{a}))[m-2])_{m-1+s} \longrightarrow \cdots\\
&\longrightarrow \textstyle\bigoplus_{\mathbf{a}\in C^{(n)}_{m-s''}(\mathbf{i})} e_{\mathbf{v^i}(\mathbf{a}')}(\tilde{P}(\mathbf{v^i}(\mathbf{a}))[m-s''])_{m+s} \longrightarrow 0,
\end{aligned}$$

with $s'' = n+1-|O(\mathbf{a}')|-|T(\mathbf{a}')|$. So we have that

$$\begin{array}{rl} & \dim{}_k e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\operatorname{Ker} f_{m-1})_{m-1+s} \\ = & \sum\limits_{t=0}^{s''}(-1)^t \dim{}_k \bigoplus_{\mathbf{a}\in C^{(n)}_{m-1-t}(\mathbf{i})} e_{\mathbf{v}^{\mathbf{i}}(\mathbf{a}')}(\tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{a}))[m-1-t])_{m-1+s} \\ = & \sum\limits_{t=0}^{s''}(-1)^t \left(\begin{array}{c} s'' \\ t \end{array}\right) = (1-1)^{s''} = 0. \end{array}$$

This proves that $(\operatorname{Ker} f_{m-1})_{m-1+s} = 0$ for $1 \le s \le n$, and thus $\operatorname{Ker} f_{m-1} = \operatorname{soc} \tilde{P}(\mathbf{v}^{\mathbf{i}}(\mathbf{b}))$. □

Let $\tilde{\Lambda}_0$ be the subspace spanned by the idempotents of $\tilde{\Lambda}(n)$, and let $\tilde{\Lambda}_1$ be the subspace spanned by the arrows. Then $\tilde{\rho}^{\bar{\mathbf{g}}}(n) \subset \tilde{\Lambda}_1 \otimes_{\tilde{\Lambda}_0} \tilde{\Lambda}_1$, and it spans a subspace $R^{\bar{\mathbf{g}}} \subset \tilde{\Lambda}_1 \otimes_{\tilde{\Lambda}_0} \tilde{\Lambda}_1$. Let $T_{\tilde{\Lambda}_0}(\tilde{\Lambda}_1) = \tilde{\Lambda}_0 + \tilde{\Lambda}_1 + \tilde{\Lambda}_1 \otimes_{\tilde{\Lambda}_0} \tilde{\Lambda}_1 + \cdots \tilde{\Lambda}_1^{\otimes_{\tilde{\Lambda}_0} t} + \cdots$ be the tensor algebra and let $(R^{\bar{\mathbf{g}}})$ be the ideal generated by $R^{\bar{\mathbf{g}}}$, then $\tilde{\Lambda}^{\bar{\mathbf{g}}}(n) \simeq T_{\tilde{\Lambda}_0}(\tilde{\Lambda}_1)/(R^{\bar{\mathbf{g}}})$ is a quadratic algebra. The quadratic dual $\tilde{\Lambda}^!(n)$ of $\tilde{\Lambda}(n)$ is defined as the quotient $T_{\tilde{\Lambda}_0}(D\tilde{\Lambda}_1)/(R^{\bar{\mathbf{g}},\perp})$, where $D\tilde{\Lambda}_1$ is the dual space of Λ_1 and $R^{\bar{\mathbf{g}},\perp}$ is the annihilator of $R^{\bar{\mathbf{g}}}$ in $D(\tilde{\Lambda}_1 \otimes_{\tilde{\Lambda}_0} \tilde{\Lambda}_1)$. From a different view point, Theorem 2.3 is restated as following theorem:

Theorem 3.4. *Assume that* $m \geq 3$.

$\tilde{\Lambda}^{\bar{\mathbf{g}}}(n)$ *is an almost Koszul algebra of type* $(n+1, m-1)$, *and its quadratical dual* $\tilde{\Lambda}^{\bar{\mathbf{g}},!}(n)$ *is almost Koszul algebra of type* $(m-1, n+1)$.

Both $\tilde{\Lambda}^{\bar{\mathbf{g}}}(n)$ *and* $\tilde{\Lambda}^{\bar{\mathbf{g}},!}(n)$ *are periodic algebras.*

More precisely, the minimal periodicity of $\tilde{\Lambda}^{\bar{\mathbf{g}}}(n)$ *is* $(n+1)m$, *and the minimal periodicity of* $\tilde{\Lambda}^{\bar{\mathbf{g}},!}(n)$ *is* $(n+1)(n+2)$.

Proof. If follows directly from Lemma 3.3 that $\tilde{\Lambda}^{\bar{\mathbf{g}}}(n)$ is an almost Koszul algebra of type $(n+1, m-1)$. So $\tilde{\Lambda}^{\bar{\mathbf{g}},!}(n)$ is almost Koszul algebra of type $(m-1, n+1)$, by Proposition 3.11 of [7].

Note that for each $\mathbf{i} \in \tilde{Q}(n)_0$, by (d) of Lemma 3.3, we have that $\Omega^m \tilde{S}(\mathbf{i}) \simeq \tilde{S}(\mathbf{v}^{\mathbf{i}}(\mathbf{b}(\mathbf{i}))) = \tilde{S}(\omega(\mathbf{i}))$. If $\mathbf{i} = (i_1, \ldots, i_n)$, then

$$\mathbf{b}(\mathbf{i}) = (m+n-1-\sum_{t=1}^{n} i_t, i_1-1, i_2-1, \ldots, i_n-1)$$

and we have

$$\omega(\mathbf{i}) = \mathbf{v}^{\mathbf{i}}(\mathbf{b}(\mathbf{i})) = (m+n-\sum_{t=1}^{n} i_t, i_1, \ldots, i_{n-1}).$$

Thus $\omega^t(\mathbf{i}) = (i_{n-t+2}, \ldots, i_n, m+n-\sum\limits_{t=1}^{n} i_t, i_1, \ldots, i_{n-t})$, and $\omega^{n+1}(\mathbf{i}) = \mathbf{i}$. This proves that $\Omega^{m(n+1)}\tilde{S}(\mathbf{i}) = \tilde{S}(\mathbf{i})$, and $n+1$ is minimal such that this holds for all $\mathbf{i} \in \tilde{Q}(n)_0$. So the minimal periodicity of $\tilde{\Lambda}^{\bar{\mathbf{g}}}(n)$ is $(n+1)m$.

Let $N(\tilde{\Lambda}^{\bar{\mathbf{g}}}(n), m-1)$ be the matrix with

$$N(\tilde{\Lambda}^{\bar{\mathbf{g}}}(n), m-1)_{\mathbf{ij}} = \dim{}_k e_{\mathbf{i}} \Omega^m \tilde{\Lambda}_0^{\bar{\mathbf{g}}}(n) e_{\mathbf{j}}.$$

$N(\tilde{\Lambda}^{\bar{\mathbf{g}}}(n), m-1)$ is the matrix of the permutation of the simples $\tilde{\Lambda}^{\bar{\mathbf{g}}}(n)$-modules defined by Ω^m, so the order of $N(\tilde{\Lambda}^{\bar{\mathbf{g}}}(n), m-1)$ is $n+1$.

Now let $N(\tilde{\Lambda}^{\bar{\mathbf{g}},!}(n), n+1)$ be the matrix with

$$N(\tilde{\Lambda}^{\bar{\mathbf{g}},!}(n), n+1)_{\mathbf{ij}} = \dim{}_k e_{\mathbf{i}} \Omega^{n+2} \tilde{\Lambda}_0^{\bar{\mathbf{g}},!}(n) e_{\mathbf{j}}.$$

$N(\tilde{\Lambda}^{\bar{\mathbf{g}},!}(n), n+1)$ is the matrix of the permutation of the simples $\tilde{\Lambda}^{\bar{\mathbf{g}}}(n), m-1)$-modules defined by Ω^{n+2}. By Proposition 3.14 of [7], the permutation matrix $N(\tilde{\Lambda}^{\bar{\mathbf{g}},!}(n), n+1)$ have the same order as $N(\tilde{\Lambda}^{\bar{\mathbf{g}}}(n), m-1)$ since they are transposes. This proves that the minimal periodicity of $\tilde{\Lambda}^{\bar{\mathbf{g}},!}(n)$ is $(n+1)(n+2)$. □

4. $(n+1)$-CUBOID TRUNCATIONS

Assume that $\Lambda(n) = \Lambda^{\mathbf{g}}(n)$ is an n-translation algebra with admissible n-translation quiver $Q(n)$. Let $\bar{\mathbf{g}} = (\mathbf{g}, g')$ be as defined in Section 2, let $\tilde{\Lambda}(n) = \tilde{\Lambda}^{\bar{\mathbf{g}}}(n)$ be a twisted trivial extension of $\Lambda(n)$. Now consider truncations on $\overline{\Lambda}(n) = \overline{\Lambda}^{\bar{\mathbf{g}}}(n) = \tilde{\Lambda}^{\bar{\mathbf{g}}}(n)\#\mathbb{Z}^*$ and we write $\overline{\rho}(n)$ for $\overline{\rho}^{\bar{\mathbf{g}}}(n)$. It follows from Theorem 2.8 that $\overline{\Lambda}(n)$ is an n-translation algebra with stable n-translation quiver $\overline{Q}(n)$ and n-translation $\overline{\tau}_n\mathbf{i} = \mathbf{i} - \mathbf{e}_{n+1}$. Write $\overline{\tau} = \overline{\tau}_n$ for the n-translation of $\overline{Q}(n)$.

Let $\overline{S}(\mathbf{i})$ be the simple $\overline{\Lambda}(n)$-module at $\mathbf{i} \in \overline{Q}(n)_0$ and let $\overline{P}(\mathbf{i})$ and $\overline{I}(\mathbf{i})$ be its projective cover and injective envelop as $\overline{\Lambda}(n)$-modules, respectively. Conventionally, set $\overline{S}(\mathbf{i})$, $\overline{P}(\mathbf{i})$ and $\overline{I}(\mathbf{i})$ to be zero when $\mathbf{i} \notin \overline{Q}(n)_0$.

For each $\mathbf{i} = (i_1, \ldots, i_n, i_{n+1}) \in \overline{Q}(n)_0$, let $\mathbf{i}' = (i_1, \ldots, i_n)$ be its n-truncation in $\tilde{Q}(n)_0 = Q(n)_0$. Define its $(n+1)$-cuboid $C^{(n)}(\mathbf{i}) = C^{(n)}(\mathbf{i}')$. We have characterization of the linear part of the projective of simple $\overline{\Lambda}^{\bar{\mathbf{g}}}(n)$-modules using the $(n+1)$-cuboids $C^{(n)}(\mathbf{i})$ similar to Proposition 3.2. Let

$$\cdots \longrightarrow \overline{P}^2(\overline{S}(\mathbf{i})) \xrightarrow{f_2} \overline{P}^1(\overline{S}(\mathbf{i})) \xrightarrow{f_1} \overline{P}^0(\overline{S}(\mathbf{i})) \xrightarrow{f_0} \overline{S}(\mathbf{i}) \longrightarrow 0 \tag{23}$$

be a minimal projective resolution of the simple $\overline{\Lambda}$-modules $\overline{S}(\mathbf{i})$ of corresponding to the vertex $\mathbf{i} \in \overline{Q}(n)_0$. If $\mathbf{i} \in \overline{Q}_0$, call $\mathbf{a}$ an $\mathbf{i}$-*quiver vertex* if $\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{a}) \in \overline{Q}(n)_0$.

Consider the linear part of this projective resolution. We have

Proposition 4.1. *For $l = 0, 1, \ldots, m-1$ and $\mathbf{i} \in \overline{Q}(n)_0$, we have that $\overline{P}^l(\overline{S}(\mathbf{i}))$ is generated in degree l and*

$$\overline{P}^l(\overline{S}(\mathbf{i})) \simeq \bigoplus_{\mathbf{a} \in C_l^{(n)}(\mathbf{i})} \overline{P}(\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{a})). \tag{24}$$

Further more, $C_{m-1}^{(n)}(\mathbf{i}) = \{\mathbf{b}\}$, and $\operatorname{Ker} f_{m-1} \simeq \operatorname{soc} \overline{P}(\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{b})) = \overline{S}(\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{b} + \mathbf{e}))$ is simple.

Recall that a full bound subquiver Q' of $\overline{Q}(n)$ is called a $\overline{\tau}_n$-*slice* of Q if it has the following property [13]:

(a) for each vertex $\mathbf{i}$ of $\overline{Q}(n)$, the intersection of the $\overline{\tau}_n$-orbit of v and the vertex set of Q' is a single-point set;

A $\overline{\tau}_n$-slice is called a *complete $\overline{\tau}_n$-slice*, if it also has the following property:

(b) Q' is path complete in the sense that for any path $p : v_0 \longrightarrow v_1 \longrightarrow \ldots \longrightarrow v_t$ of $\overline{Q}(n)$ with v_0 and v_t in Q', the whole path p lies in Q'.

The algebra defined by a complete $\overline{\tau}$-slice of $\overline{Q}(n)$ is called a $\overline{\tau}$-*slice algebra* of $\overline{Q}(n)$ (or of $\overline{\Lambda}(n)$).

Let $Q(n) \times \{t\}$ be the full bound subquiver of $\overline{Q}(n)$ with vertex set $Q(n)_0 \times \{t\}$ for any $t \in \mathbb{Z}$. It is easy to check that we have the following result.

Proposition 4.2. *$Q(n) \times \{t\}$ are complete $\overline{\tau}$-slices of $\overline{Q}(n)$ for all t.*

Let $\Lambda(n,t)$ be the algebra defined by the bound quiver $Q(n)\times\{t\}$. $\Lambda(n,t)$ are isomorphic to $\Lambda^{\mathbf{g}}(n)$ for all t.

Now we define the $(n+1)$-*cuboid completion* $Q(n+1)(t)$ of $Q(n)\times\{t\}$ as the full subquiver of $\overline{Q}(n)$ with the vertex set:

$$Q(n+1)(t)_0 = \cup_{\mathbf{i}\in Q(n)_0\times\{t\}}\{\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{a})|\mathbf{a}\in C^{(n)}(\mathbf{i})\}.$$

Clearly, $Q(n+1)(t)$ are isomorphic quivers for all t.

Let $Q(n+1)(1)^{(t)}$ and $Q(n+1)^{(t)}$ be the full subquivers of $Q(n+1)(1)$ and $Q(n+1)$ with vertex set $Q(n+1)(1)^{(t)}_0 = \{\mathbf{i}=(i_1,\dots,i_t,1,\dots,1)\in Q(n+1)(1)_0\}$ and $Q(n+1)^{(t)}_0 = \{\mathbf{i}=(i_1,\dots,i_t,1,\dots,1)\in Q(n+1)_0\}$, respectively.

Lemma 4.3. *For $1\le t\le n+1$, $(Q(t),\rho(t))$ can be identify with*

$$(Q(n+1)(1)^{(t)},\overline{\rho}(n)[Q(n+1)(1)^{(t)}])$$

as full bound subquivers.

Proof. We prove by using induction on n and t. The assertion clearly holds for $n=0$ since $Q(1)(1)^{(1)}=Q(1)$.

Assume that $n>0$ and the lemma holds for $n'<n$. By inductive assumption $(Q(t),\rho(t))$ can be identify $Q(n)(1)^{(t)}$, and $Q(n)(1)=Q(n)(1)^{(n)}$, which is identified with $Q(n)$ and with $Q(n)\times\{1\}=Q(n+1)(1)^{(n)}$, a full subquiver of $Q(n+1)$ inside $Q(n+1)(1)$, if $t\le n$.

We need only to prove the case of $t=n+1$.

For any $\mathbf{i}=(i_1,\dots,i_n,i_{n+1})\in Q(n+1)_0$, we have that $(i_1,\dots,i_n)\in Q(n)_0$ and $i_{n+1}\le m+n-1-\sum_{t=1}^{n} i_t$, by definition. It follows easily that $i_n+i_{n+1}-1\le m+n-1-\sum_{t=1}^{n-1} i_t$, thus $\mathbf{i}'=(i_1,\dots,i_{n-1},i'_n=i_n+i_{n+1}-1,1)\in Q(n)_0\times\{1\}$. Let $\mathbf{a}=(0,\dots,0,i_{n+1}-1)$, then $\mathbf{a}\in C^{(n)}(\mathbf{i}')$, and we have that

$$\bar{\mathbf{v}}^{\mathbf{i}'}(\mathbf{a})=(i_1,\dots,i_n,i_{n+1})=\mathbf{i}.$$

This proves that $Q(n+1)_0\subset Q(n+1)(1)_0$.

On the other hand, for any $\mathbf{i}=(i_1,\dots,i_n,i_{n+1})\in Q(n+1)(1)$, there is an $\mathbf{i}'=(i'_1,\dots,i'_n,1)\in Q(n+1)_0\times\{1\}$ such that $\mathbf{i}=\bar{\mathbf{v}}^{\mathbf{i}'}(\mathbf{a})$ for some $\mathbf{a}\in C^{(n)}(\mathbf{i}')$. Thus $i_t=i'_t+a_t-a_{t+1}>1$ for $t\le n$ and $i_{n+1}=1+a_{n+1}$. For $1\le s\le n$, we have

$$\begin{aligned}\sum_{t=1}^{s} i_t &= \textstyle\sum_{t=1}^{s} i'_t+a_1-a_{s+1}\\ &\le \textstyle\sum_{t=1}^{s} i'_t+m+n-1-\displaystyle\sum_{t=1}^{n} i'_t-a_{s+1}\\ &= m+n-1-\sum_{t=s+1}^{n} i'_t-a_{s+1}\le m+s-1,\end{aligned}$$

and

$$\sum_{t=1}^{n+1} i_t=\sum_{t=1}^{n+1} i'_t+a_1\le m+n.$$

This proves that $\mathbf{i}\in Q(n+1)_0$ and thus $Q(n+1)_0=Q(n+1)(1)_0$.

Now we prove that $Q(n+1)(1)$ is a full bound subquiver of $\overline{Q}(n)$. Assume that $\mathbf{j},\mathbf{j}(t),\mathbf{j}(s,t)\in Q(n+1)(1)_0$ and $\mathbf{j}(s)\in\overline{Q}(n)_0$. Then $\mathbf{i}=\mathbf{j}-(j_{n+1}-1)\mathbf{e}_{n+1}\in Q(n+1)(1)^{(n)}_0$ and $\mathbf{i}(s)=\mathbf{j}(s)-(j_{n+1}-1)\mathbf{e}_{n+1},\mathbf{i}(t)=\mathbf{j}(t)-(j_{n+1}-1)\mathbf{e}_{n+1},\mathbf{i}(s,t)=\mathbf{j}(s,t)-(j_{n+1}-1)\mathbf{e}_{n+1}\in Q(n+1)(1)^{(n)}_0$. Thus $\mathbf{e}_t+(j_{n+1}-1)\mathbf{e}_{n+1},\mathbf{e}_t+\mathbf{e}_s+(j_{n+1}-1)\mathbf{e}_{n+1}\in C^{(n)}(\mathbf{i})$, so $\mathbf{e}_s+(j_{n+1}-1)\mathbf{e}_{n+1}\in C^{(n)}(\mathbf{i})$ and $\mathbf{j}(s)=\mathbf{v}^{\mathbf{i}}(\mathbf{e}_s+(j_{n+1}-1)\mathbf{e}_{n+1})\in Q(n+1)(1)_0$. This proves that $(Q(n+1)(1),\overline{\rho}(n)[Q(n+1)(1)])$ is a full bound quiver of $(\overline{Q}(n), \overline{\rho}(n))$, since elements of $\overline{\rho}(n)$ are either paths of length 2, or linear combinations of two paths of length 2 of the arrows of the same types. □

Write $\rho(n+1)(1)^{(t)} = \overline{\rho}(n)[Q(n+1)(1)^{(t)}]$, and $\rho(n+1)(t) = \overline{\rho}(n)[Q(n+1)(t)]$. As a corollary of Lemma 4.3, we have

Corollary 4.4. *$(Q(n+1)(t), \rho(n+1)(t))$ are full bound subquivers of $(\overline{Q}(n), \overline{\rho}(n))$.*

Let $\Lambda(n+1)(t)$ be the algebra defined by the bound quiver $(Q(n+1)(t), \rho(n+1)(t))$ and they will be called $(n+1)$-*cuboid truncations* of $\overline{\Lambda}^{\mathbf{g}}(n)$.

We have the following theorem.

Theorem 4.5. (a) *$\Lambda(n+1)(t)$ are all isomorphic to $\Lambda(n+1)(1) = \Lambda(n+1)$.*
- (b) *Let $I(t)$ be the ideal of $\overline{\Lambda}(n)$ generated by the set $\{e_{\mathbf{i}} | \mathbf{i} \in \overline{Q}(n)_0 \backslash Q(n+1)(t)_0\}$, then $\Lambda(n+1) \simeq \overline{Q}(n)/I(t)$ for all t.*
- (c) *$\Lambda(n+1)$ is n-translation algebra.*

Proof. (a), (b) follows directly from Lemma 4.3.

Now we prove (c). Identify $Q(n+1)$ with $Q(n+1)(1)$. As a truncation of an n-translation quiver, it follows from (a) and (b) that $Q(n+1)$ is an n-translation quiver.

For each $\mathbf{i} \in Q(n+1)_0 = Q(n+1)(1)_0 \subset \overline{Q}(n)_0$, we have a minimal projective resolution (23) of simple $\overline{\Lambda}(n)$-module $\overline{S}(\mathbf{i})$. Note that $\Lambda(n+1) \simeq \overline{\Lambda}(n)/I$, where I is the ideal generated by $\{e_{\mathbf{j}} | \mathbf{j} \in \overline{Q}(n)_0 \setminus Q(n+1)(1)_0\}$. Since $Q(n+1)(1)_0$ is finite let $e = \sum_{\mathbf{i} \in Q(n+1)(1)_0} e_{\mathbf{i}}$, then $\bar{e}$ is the unit of $\overline{\Lambda}(n)/I$. We have $S(\mathbf{i}) = \overline{S}(\mathbf{i})$ for $\mathbf{i} \in Q(n+1)(1)_0$. Tensor (23) with $\overline{\Lambda}(n)/I$, we get a complexes of $\Lambda(n+1)$-modules

$$\begin{aligned} \cdots \longrightarrow \overline{\Lambda}(n)/I \otimes_{\overline{\Lambda}(n)} \overline{P}^2(\overline{S}(\mathbf{i})) &\xrightarrow{1 \otimes f_2} \overline{\Lambda}(n)/I \otimes_{\overline{\Lambda}(n)} \overline{P}^1(\overline{S}(\mathbf{i})) \\ \xrightarrow{1 \otimes f_1} \overline{\Lambda}(n)/I \otimes_{\overline{\Lambda}(n)} \overline{P}^0(\overline{S}(\mathbf{i})) &\xrightarrow{1 \otimes f_0} \overline{\Lambda}(n)/I \otimes_{\overline{\Lambda}(n)} \overline{S}(\mathbf{i}) = S(\mathbf{i}) \longrightarrow 0. \end{aligned} \tag{25}$$

Note that $\overline{\Lambda}(n)/I \otimes_{\overline{\Lambda}(n)} \overline{P}^t(\overline{S}(\mathbf{i})) \simeq \overline{P}^t(\overline{S}(\mathbf{i}))/I\overline{P}^t(\overline{S}(\mathbf{i}))$. For any $y = \sum_p \bar{a}_p \otimes y_p = \bar{e} \otimes \sum_p a_p y_p$ in $\operatorname{Ker}(1 \otimes f_t)$, we have that $\sum_p a_p y_p \in \operatorname{Ker} f_t = f_{t+1}(\overline{P}^{t+1}(\overline{S}(\mathbf{i})))$ and hence there is an $x \in \overline{P}^{t+1}(\overline{S}(\mathbf{i}))$ such that $f_{t+1}(x) = \sum_p a_p y_p$. Thus $(1 \otimes f_{t+1})(\bar{e} \otimes x) = y$. This proves that (25) is exact and hence a projective resolution $S(\mathbf{i})$ as $\Lambda(n+1)$-modules.

Note that $\overline{\Lambda}(n)/I \otimes_{\overline{\Lambda}(n)} \overline{P}^t(\overline{S}(\mathbf{i})) \simeq \overline{P}^t(\overline{S}(\mathbf{i}))/I\overline{P}^t(\overline{S}(\mathbf{i}))$. Thus if $\mathbf{i} \in \overline{Q}(n)_0 \setminus Q(n+1)(1)_0$, then $\overline{\Lambda}(n)/I \otimes_{\overline{\Lambda}(n)} \overline{\Lambda}(n)e_{\mathbf{i}} = (\overline{\Lambda}(n)/I\overline{\Lambda}(n))e_{\mathbf{i}} \otimes_{\overline{\Lambda}(n)} e_{\mathbf{i}} = 0$ and if $\mathbf{i} \in Q(n+1)(1)_0$, then $\overline{\Lambda}(n)/I \otimes_{\overline{\Lambda}(n)} \overline{\Lambda}(n)e_{\mathbf{i}} = (\overline{\Lambda}(n)/I\overline{\Lambda}(n)e_{\mathbf{i}} \simeq \Lambda(n+1)e_{\mathbf{i}}$. If $\mathbf{i} \in Q(n+1)(1)_0$, by Proposition 4.1, we have for $l = 0, 1, \ldots, m-1$,

$$\overline{\Lambda}(n)/I \otimes_{\overline{\Lambda}(n)} \overline{P}^l(\overline{S}(\mathbf{i})) \simeq \bigoplus_{\mathbf{a} \in C_l^{(n)}(\mathbf{i}), \mathbf{v}^{\mathbf{i}}\mathbf{a} \in Q(n+1)(1)_0} \Lambda(n+1)(1)e_{\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{a})} \tag{26}$$

is generated in degree l and $\operatorname{Ker}(1 \otimes f_{m-1})$ is either simple or zero.

This proves that $\Lambda(n+1)$ is almost Koszul of type $(n+1, m-1)$ and thus is n-translation algebra.

□

We call $\Lambda(n+1)$ a $(n+1)$-*cuboid completion of* $\Lambda(n)$.

5. Pyramid n-cubic Algebras and n-almost Split Sequences

Now fix $m \geq 3$, and a integer $n \geq 1$. Let $Q(n)$ and $\rho(n) = \rho^{\mathbf{g}}(n)$ be the data for a pyramid shaped n-cubic quiver $Q(n) = Q^{\mathbf{g}}(n)$ as defined in (2), (3) and (4), and let $\Lambda(n)$ be the pyramid n-cubic algebra defined by the bound quiver $(Q(n), \rho(n))$.

For a bound path p from i to j such that any of bound path from i to j of the same length as p is linearly dependent, recall that p is *left stark of degree* t with respect to i' if pw is a bound element for any bound element w from i' to i of length $t < n + 1 - l(p)$; and p is *right stark of degree* t with respect to j' if wp is a bound element for any bound element w' from j to j' of length $t < n + 1 - l(p)$. A bound path p is *shiftable* if it is linear dependent to paths of the form $p'p''$ with p'' is trivial or linearly dependent on paths passing through no injective vertex and p' is trivial or linearly dependent on paths passing through no projective vertex.

Recall that an n-translation quiver Q is called *admissible* if it satisfies the following conditions:

(i) For each bound path p, there are paths q' and q'' such that $q'pq''$ is a bound path of length $n + 1$.
(ii) Any bound path p from a non-injective vertex i to a non-projective vertex j is linearly dependent to shiftable paths.
(iii) Let i be a non-projective vertex. Let p be a bound path ending at i and let q be a bound path starting at τi with $l(p) + l(q) \leq n$. If p passes a projective vertex and q passes an injective vertex, then p is either left stark with respect to $t(q)$, or q is right stark with respect to $s(p)$, of length $n + 1 - (l(p) + l(q))$.

We have the following lemma describing the bound paths in $Q(n)$

Lemma 5.1. *Let* $q = \gamma^{(t_r)}_{\mathbf{i}(t_1,\cdots,t_{r-1})}\gamma^{(t_{r-1})}_{\mathbf{i}(t_1,\cdots,t_{r-2})}\cdots\gamma^{(t_1)}_{\mathbf{i}}$ *be a bound path in* $Q(n)$.

If $t_r \neq 1$ *and* $i_{t_r-1} > 1$, *then there is a* $d \in k$ *such that*

$$q = d\gamma^{(t_{r-1})}_{\mathbf{i}(t_1,\cdots,t_{r-2},t_r)}\gamma^{(t_{r-2})}_{\mathbf{i}(t_1,\cdots,t_{r-3},t_r)}\cdots\gamma^{(t_1)}_{\mathbf{i}(t_r)}\gamma^{(t_r)}_{\mathbf{i}}.$$

If $j_{t_1} > 1$, *then there is a* $d \in k$ *such that*

$$q = d\gamma^{(t_1)}_{\mathbf{i}(t_2,\cdots,t_{r-1},t_r)}\gamma^{(t_r)}_{\mathbf{i}(t_2,\cdots,t_{r-1})}\cdots\gamma^{(t_3)}_{\mathbf{i}(t_2)}\gamma^{(t_2)}_{\mathbf{i}}.$$

Proof. Set $\mathbf{j}^{(u)} = \mathbf{i}(t_1, \ldots, t_{u-1}) = (j^{(u)}_1, \ldots, j^{(u)}_{n+1})$ for $u = 1, \ldots, r$. Since q is a bound path, $t_1 \neq t_r$ are pairwise different.

If $t_r \neq 1$ and $i_{t_r-1} > 1$, then for $u = 1, \ldots, r-1$, $j^{(u)}_{t_r-1} \geq i_{t_r-1} > 1$, and we have that $\mathbf{j}^{(u)}(t_r) = \mathbf{j}^u - \mathbf{e}_{t_r-1} + \mathbf{e}_{t_r}$ is a vertex in $Q(n)$, too. Thus there are nonzero $d_1, \ldots, d_{r-1} \in k$ such that $\gamma^{(t_r)}_{\mathbf{i}(t_1,\ldots,t_s)}\gamma^{(t_s)}_{\mathbf{i}(t_1,\ldots,t_{s-1})} = d_s\gamma^{(t_s)}_{\mathbf{i}(t_1,\ldots,t_{s-1},t_r)}\gamma^{(t_r)}_{\mathbf{i}(t_1,\ldots,t_{s-1})}$ and thus

$$\begin{array}{rcl} q & = & \gamma^{(t_r)}_{\mathbf{i}(t_1,\cdots,t_{r-1})}\gamma^{(t_{r-1})}_{\mathbf{i}(t_1,\cdots,t_{r-2})}\cdots\gamma^{(t_2)}_{\mathbf{i}(t_1)}\gamma^{(t_1)}_{\mathbf{i}} \\ & = & d_{r-1}\gamma^{(t_{r-1})}_{\mathbf{i}(t_1,\cdots,t_{r-2},t_r)}\gamma^{(t_r)}_{\mathbf{i}(t_1,\cdots,t_{r-2})}\cdots\gamma^{(t_2)}_{\mathbf{i}(t_1)}\gamma^{(t_1)}_{\mathbf{i}} \\ & = & \cdots \\ & = & d_{r-1}\cdots d_1\gamma^{(t_{r-1})}_{\mathbf{i}(t_1,\cdots,t_{r-2},t_r)}\gamma^{(t_{r-2})}_{\mathbf{i}(t_1,\cdots,t_{r-3},t_r)}\cdots\gamma^{(t_1)}_{\mathbf{i}(t_r)}\gamma^{(t_r)}_{\mathbf{i}} \\ & = & d\gamma^{(t_{r-1})}_{\mathbf{i}(t_1,\cdots,t_{r-2},t_r)}\gamma^{(t_{r-2})}_{\mathbf{i}(t_1,\cdots,t_{r-3},t_r)}\cdots\gamma^{(t_1)}_{\mathbf{i}(t_r)}\gamma^{(t_r)}_{\mathbf{i}}. \end{array}$$

Similarly, if $t_1 > 1$, then we have $j^{(u)}_{t_1-1} > 1$ for $u = 2, \ldots, n$ so $(t_1)\mathbf{j}^{(u)} = \mathbf{j}^{(u)} + \mathbf{e}_{t_r-1} - \mathbf{e}_{t_r} \in Q(n)_0$. This implies $(t_1)\mathbf{j}^{(u)} \in Q(n+1)_0$ and there are nonzero

$d'_2, \dots, d'_r \in k$ such that $\gamma^{(t_s)}_{\mathbf{i}(t_1,\dots,t_{s-1})}\gamma^{(t_1)}_{\mathbf{i}(t_2,\dots,t_{s-1})} = d'_s\gamma^{(t_1)}_{\mathbf{i}(t_2,\dots,t_s)}\gamma^{(t_s)}_{\mathbf{i}(t_2,\dots,t_{s-1})}$ and thus

$$\begin{array}{rcl} q & = & \gamma^{(t_r)}_{\mathbf{i}(t_1,\cdots,t_{r-1})}\gamma^{(t_{r-1})}_{\mathbf{i}(t_1,\cdots,t_{r-2})}\cdots\gamma^{(t_2)}_{\mathbf{i}(t_1)}\gamma^{(t_1)}_{\mathbf{i}} \\ & = & d'_2\gamma^{(t_r)}_{\mathbf{i}(t_1,\cdots,t_{r-1})}\gamma^{(t_{r-1})}_{\mathbf{i}(t_2,\cdots,t_{r-2})}\cdots\gamma^{(t_1)}_{\mathbf{i}(t_2)}\gamma^{(t_2)}_{\mathbf{i}} \\ & = & \cdots \\ & = & d'_r d'_{r-1}\cdots d'_2\gamma^{(t_1)}_{\mathbf{i}(t_2,\cdots,t_r)}\gamma^{(t_r)}_{\mathbf{i}(t_2,\cdots,t_{r-1})}\cdots\gamma^{(t_3)}_{\mathbf{i}(t_2)}\gamma^{(t_2)}_{\mathbf{i}} \\ & = & d'\gamma^{(t_1)}_{\mathbf{i}(t_2,\cdots,t_r)}\gamma^{(t_r)}_{\mathbf{i}(t_2,\cdots,t_{r-1})}\cdots\gamma^{(t_3)}_{\mathbf{i}(t_2)}\gamma^{(t_2)}_{\mathbf{i}}. \end{array}$$

□

Theorem 5.2. *Pyramid n-cubic algebra $\Lambda(n)$ is an extendible $(n-1)$-translation algebra with admissible $(n-1)$-translation quiver $Q(n)$.*

Proof. We prove theorem by using induction on n. We have

$$Q(1) = Q : \overset{1}{\circ}\longrightarrow\overset{2}{\circ}\longrightarrow\circ\cdots\circ\longrightarrow\overset{m}{\circ}, \tag{27}$$

with the relation $\rho(1) = \{\alpha_{i+1}\alpha_i | i = 1, \dots, m-2\}$. Clearly, $Q(1) = (Q(1), \rho(1))$ is a admissible 0-translation quiver with 0-translation $\tau_0 : i+1 \to i$ for $i = 1, \dots, m-1$. $\Lambda(1)$ is a Koszul algebra with radical squared zero. So it is a 0-translation algebra, and it is extendable by Theorem 2.1 of [7].

Assume that $\Lambda(n)$ is an extendible $(n-1)$-translation algebra and its bound quiver $Q(n)$ is an admissible $(n-1)$-translation quiver.

It follows from Theorem 4.5 that $\Lambda(n+1)$ is an n-translation algebra with n-translation $Q(n+1)$ and n-translation $\tau_n : \mathbf{i} \to \mathbf{i} - \mathbf{e}_{n+1}$. By Theorem 3.4, $\Lambda(n+1)$ is extendible. We need only to prove that $Q(n+1)$ is admissible.

Let $\mathcal{P}$ and $\mathcal{I}$ be the sets of projective vertices and injective vertices of $Q(n+1)$, respectively. Then

$$\mathcal{P} = \{\mathbf{i} \in Q(n+1)_0 | i_{n+1} = 1\} \qquad \mathcal{I} = \{\mathbf{i} \in Q(n+1)_0 | \, |\mathbf{i}| = m+n\}.$$

Let $p = \gamma^{(t_s)}_{\mathbf{i}(t_1,\cdots,t_{s-1})}\cdots\gamma^{(t_1)}_{\mathbf{i}}$ be a bound path in $\Lambda(n+1)$. If $t_h = 1$, then

$$\mathbf{i}, \mathbf{i}(t_1), \dots, \mathbf{i}(t_1, \dots, t_{h-1})$$

are not injective. If $t_h = n+1$, then

$$\mathbf{i}(t_1, \dots, t_h), \mathbf{i}(t_1 \dots, t_h, t_{h+1}), \dots, \mathbf{i}(t_1, \dots, t_s)$$

are not projective.

Assume that p is a bound path from $\mathbf{i}$ to $\mathbf{j}$ in $Q(n+1)$. Then $\mathbf{j} \in \{\mathbf{v}^{\mathbf{i}}(\mathbf{a}) | \mathbf{a} \in U^{n+1}\}$ and $\mathbf{i} \in \{\mathbf{v}^{\mathbf{j}-\mathbf{e}_{n+1}}(\mathbf{a}) | \mathbf{a} \in U^{n+1}\}$. If $\mathbf{j}$ is non-projective, then there is a path p' from $\tau_n\mathbf{j}$ to $\mathbf{i}$ such that pp' is a bound path of length $n+1$, take $p'' = e_{\mathbf{j}}$ be the trivial path. If $\mathbf{i}$ is non-injective then there is a path p'' from $\mathbf{j}$ to $\tau_n^{-1}\mathbf{i}$ such that $p''p$ is a bound path of length $n+1$, take $p' = e_{\mathbf{i}}$ be the trivial path. If $\mathbf{i}$ is injective and $\mathbf{j}$ is projective, then we have that $\sum\limits_{t=1}^{n+1} i_t = m+n$ and $j_{n+1} = 1$. So $i_{n+1} = 1$ and $|\mathbf{j}| = m+n$, and all the vertices on the path p are projective and injective, which is on the subquiver $Q(n) \times \{1\}$ of $Q(n+1)$. By the inductive assumption, there are paths q' and p'' in $Q(n) \times \{1\}$, such $q'pp''$ is a bound path from $\mathbf{i}' = (i'_1, \dots, i'_{n+1})$ to $\mathbf{j}'(j'_1, \dots, j'_{n+1})$ of length n. But $i'_t \geq 1$ for $t = 1, \dots, n+1$ and $j'_t \geq 1$, $q'pq''$ is formed by arrows of different type, thus $j'_n > 1$ and $\mathbf{j}', \mathbf{j}'(n+1)$ are vertices in $Q(n+1)$. So $\gamma^{(n+1)}_{\mathbf{j}'} : \mathbf{j}' \to \mathbf{j}'(n+1)$ is an arrow of $Q(n+1)$. Let $p' = \gamma^{(n+1)}_{\mathbf{j}'}q'$, then $p'pp''$ is a bound path of length $n+1$ in $Q(n+1)$. This proves the Condition (i) of the admissibility.

Let $p = \gamma^{(t_s)}_{\mathbf{i}(t_1,\cdots,t_{s-1})} \cdots \gamma^{(t_1)}_{\mathbf{i}}$ be a bound path from non-injective vertex $\mathbf{i}$ to non-projective vertex $\mathbf{j}$. Then we have that $j_{n+1} > 1$ and $|\mathbf{i}| < m+n$. We prove that p is shiftable by using induction on s.

If $s = 1$, then either $t_1 = 1$, then $i_{n+1} = j_{n+1}$, both $\mathbf{i}$ and $\mathbf{j}$ are non-projective and we take $p' = p$, $p'' = e_{\mathbf{i}}$, so $p = p'p''$ is shiftable, or $t_1 \neq 1$ and $|\mathbf{j}| = |\mathbf{i}| < m+n$, thus both $\mathbf{i}$ and $\mathbf{j}$ are non-injective and we take $p' = e_{\mathbf{j}}$, $p'' = p$, so $p = p'p''$ is shiftable p.

Assume that $l \geq 1$ is an integer and any bound path of length l from a non-injective vertex to a non projective vertex is shiftable for $s = l$.

Now let $s = l+1$. Assume that p passes through both projective and injective vertex. Since for any arrow $\alpha : \mathbf{i}' \to \mathbf{j}'$, $\mathbf{j}'$ is injective if $\mathbf{i}'$ is so, and $\mathbf{i}'$ is projective if $\mathbf{j}'$ is so, thus $\mathbf{i}$ is projective and $\mathbf{j}$ is injective and we have $i_{n+1} = 1$ and $|\mathbf{j}| = m+n$. This implies that there is one arrow of type 1 one arrow of type $n+1$ in p in the path, and we have $j_{n+1} = 2$ and $|\mathbf{i}| = m+n-1$.

If $t_1 \neq 1$, p does not start with an arrow of type 1, then $\mathbf{i}(t_1)$ is not injective, then $p' = \gamma^{(t_s)}_{\mathbf{i}(t_1,\cdots,t_{s-1})} \cdots \gamma^{(t_2)}_{\mathbf{i}(t_1)}$ is a bound path from a non-injective vertex $\mathbf{i}(t_1)$ to a non-projective vertex $\mathbf{j}$ and thus it is shiftable by inductive assumption. So p is also shiftable by definition.

If $t_l \neq n+1$, that is, p does not end with an arrow of type $n+1$, then $(t_l)\mathbf{j}$ is not projective, then $p' = \gamma^{(t_l)}_{\mathbf{i}(t_1,\cdots,t_{l-1})} \cdots \gamma^{(t_1)}_{\mathbf{i}}$ is a bound path from a non-injective vertex $\mathbf{i}$ to a non-projective vertex $(t_l)\mathbf{j}$ and thus it is shiftable by inductive assumption. So p is also shiftable by definition.

Now assume that $t_1 = 1$, $t_{l+1} = n+1$.

If $i_{t_r-1} = 1$ for $1 < r \leq l+1$ and $j_{t_r} = 1$ for $1 \leq r \leq l$. Then for $r = 1, \ldots, l+1$, $\mathbf{i}(t_1, \ldots, t_r) \in Q(n+1)$ if and only if $t_v = v$ for $v = 1, \ldots, r$ and similarly, for $r = 1, \ldots, l$, $(t_r, \ldots, t_{l+1})\mathbf{j} \in Q(n+1)$ if and only if $t_{l+1-v} = n+1-v$ for $v = 1, \ldots, l+1-r$. This implies $l = n$ and $j_{n+1} = i_{n+1} + 1 = m$, since $\mathbf{j}$ is injective and $|\mathbf{j}| = m+n$. So $i_{n+1} = m-1 \geq 2$, this contradicts the fact that $\mathbf{i}$ is projective.

So there is $1 < r \leq l+1$ such that $i_{t_r-1} > 1$, or there is $1 \leq r \leq l$, such that $j_{t_r} > 1$. In the first case we have that

$$p = d\gamma^{(t_{l+1})}_{\mathbf{i}(t_1,\cdots,t_l)} \cdots \gamma^{(t_{r+1})}_{\mathbf{i}(t_1,\cdots,t_r)} \gamma^{(t_{r-1})}_{\mathbf{i}(t_1,\cdots,t_{r-2},t_r)} \cdots \gamma^{(t_1)}_{\mathbf{i}(t_r)} \gamma^{(t_r)}_{\mathbf{i}}$$

by lemma 5.1, and p is a multiple of a path which does not start with an arrow of type 1. Similarly, in the second we have that

$$p = d\gamma^{(t_r)}_{\mathbf{i}(t_1,\ldots,t_{r-1},t_{r+1},\ldots,t_{l+1})} \gamma^{(t_{l+1})}_{\mathbf{i}(t_1,\cdots,t_{r-1},t_{r+1},\ldots,t_l)} \cdots \gamma^{(t_{r+1})}_{\mathbf{i}(t_1,\cdots,t_{r-1})} \gamma^{(t_{r-1})}_{\mathbf{i}(t_1,\cdots,t_{r-2})} \cdots \gamma^{(t_1)}_{\mathbf{i}(t_r)}.$$

Thus p is linearly dependent to shiftable paths by above argument and (ii) of the admissible condition hold.

Since for any pair of vertex $\mathbf{i}, \mathbf{j}$ of $Q(n+1)$, all the bound paths from i to j are linearly dependent, if there is one, it is easy to see that (iii) of the admissible condition holds.

This proves that a pyramid n-cubic algebra is $(n-1)$-translation algebra with admissible $(n-1)$-translation quiver, by induction. $\square$

Let $\Gamma(n+1) = \Lambda(n+1)^{!^{op}}$ be the Koszul dual of $\Lambda(n+1)$. They have the same quiver with quadratic dual relations. The Koszul complexes gives a correspondence between the radical layers of the projective cover of a simple and terms in the projective resolution of the corresponding simple of these two algebras [7]. The the radical layers of the projective cover and the terms of projective resolution of a simple $\Lambda(n)$-module are described by the n-cubic cells $H^{\mathbf{i}}$ and n-hammocks $H^{\mathbf{i}}_C$ (see Lemma 2.7 and (26)).

Now we consider the n-almost split sequence in the category of the finite generated projective modules. We first study the n-cubic cells and n-hammocks, we have the following lemma.

Lemma 5.3. *Let $Q(n)$ be a pyramid shaped n-cubic quiver and let $\mathbf{i}$ be a vertex of $Q(n)$.*

If $H^{\mathbf{i}}$ is complete, then $H_C^{\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{e})}$ is not complete.

If $H^{\mathbf{i}}_C$ is complete, then $H^{\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{b}(\mathbf{i}))}$ is not complete.

Proof. For $\mathbf{i} = (i_1, \dots, i_n) \in G(n)_0$, then $\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{e}) = (i_1, \dots, i_n + 1)$. Thus $\mathbf{b}(\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{e})) = (m+n-1-\sum_{t=1}^n i_t, i_2 - 1, \dots, i_{n-1} - 1, i_n)$ and

$$\bar{\mathbf{v}}^{\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{e})}(\mathbf{b}(\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{e}))) = (i_1+m+n-1-\sum_{t=1}^{n} i_t - i_2 + 1, 2i_2 - i_3, \dots, 2i_{n-1} - i_n - 1, 2i_n + 1),$$

and we have

$$|\bar{\mathbf{v}}^{\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{e})}(\mathbf{b}(\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{e})))| = m + n - 1 + 1 = m + n.$$

This implies that $\bar{\mathbf{v}}^{\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{e})}(\mathbf{b}(\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{e}))) \notin Q(n)_0$ and $H_C^{\bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{e})}$ is not complete.

The other statement is proven similarly. ☐

The following Lemma follows directly.

Lemma 5.4. *The following are equivalent for $\Lambda(n+1)$.*

- (i) *$\Lambda(n+1)e_{\mathbf{i}}$ is projective injective.*
- (ii) *The Loewy length of $\Lambda(n+1)e_{\mathbf{i}}$ is $n+2$.*
- (iii) *$H^{\mathbf{i}}$ is complete.*

It follows from Theorem 6.2 of [15], that $\tilde{\Lambda}(n)^{!^{op}}$ and $\overline{\Lambda}(n)^{!^{op}}$ are self-injective of Loewy length m, so they are $(m-2)$-translation algebras. Via the Koszul complexes, $(n+1)$-hammocks can be used to describe the radical layers of the projective covers and $(n+1)$-cubic cells can be used to describe the projective resolutions of the simple $\overline{\Lambda}(n+1)^{!^{op}}$-modules. As a truncation of $\overline{\Lambda}(n+1)^{!^{op}}$ we also have the following version of Lemma 5.4 for $\Gamma(n+1)$.

Lemma 5.5. *The following are equivalent for $\Gamma(n+1)$.*

- (i) *$\Gamma(n+1)e_{\mathbf{i}}$ is projective injective.*
- (ii) *The Loewy length of $\Gamma(n+1)e_{\mathbf{i}}$ is m.*
- (iii) *$H^{\mathbf{i}}_C$ is complete.*

Then we have the following result:

Theorem 5.6. *$\Gamma(n+1)$ is an $(m-2)$-translation algebra.*

$\operatorname{add}\Gamma(n+1)$ has n-almost split sequence.

$\operatorname{add}\Lambda(n+1)$ has $(m-3)$-almost split sequence.

Proof. Similar to the case for $\Lambda(n+1)$, it is easy to see that

$$\Gamma(n+1) \simeq \overline{\Gamma}(n)/I',$$

where I' is the ideal of $\overline{\Gamma}(n)$ generated by the set $\{e_{\mathbf{j}}|\mathbf{j} \in \overline{Q}(n)_0 \setminus Q(n+1)_0\}$ of idempotents. Thus $\Gamma(n+1)$ is an $(n+1)$-translation algebra.

Note that the n-translation of $\Lambda(n+1)$ is defined by $\tau_n^{-1}\mathbf{i} = \mathbf{i} + \mathbf{e}$ and the $m-2$-translation of $\Gamma(n+1)$ is defined by $\tau_{[m-2]}^{-1}\mathbf{i} = \bar{\mathbf{v}}^{\mathbf{i}}(\mathbf{b}(\mathbf{i}))$. The last two assertion follows from Theorem 7.2 of [15], and our Lemma 5.4. $\square$

We have the following version of Iyama's cone construction of absolutely $(n+1)$-complete algebras from an absolutely n-complete algebra.

Theorem 5.7. *Let $\Lambda^{\mathbf{g}}(n) = T_m^{(n)^!}(k)$ be the pyramid n-cubic algebra defined in Proposition 2.2, then there is an $(n+1)$-cuboid completion $\Lambda^{\bar{\mathbf{g}}}(n+1)$ of $\Lambda^{\mathbf{g}}(n)$ such that $T_m^{(n+1)}(k) \simeq \Lambda^{\bar{\mathbf{g}}^!}(n+1)$.*

References

[1] Assem I, Skowronski A, Simson D. Elements of the Representation Theory of Associative Algebras, I. London Mathematical Society Texts 65()

[2] Aspinwall P S, Fidkowski L M. Superpotentials for Quiver Gauge Theories, DUKE-CGTP-05-05, SLAC-PUB-11261, SU-ITP-05/21. JHEP 0610 (2006) 047.

[3] Auslander, M., Rational singularities and almost spliting sequences. Trans. Amer. Math. Soc. 293 511-531(1986)

[4] Auslander, M., Reiten, I., and Smalø, S.: *Representation theory of artin algebras*, Cambridge Studies in Advanced Math. 36, Cambridge Univ. Press(1995)

[5] Buan, A., Iyama,O., Reiten,I., Scott, J.: Cluster structures for 2-Calabi-Yau categories and unipotent groups, Compos. Math. 145 1035-1079(2009)

[6] Baer, D., Geigle, W., Lenzing, H.: The preprojective algebra of a tame hereditary Artin algebra, Comm. Algebra 15 425-457(1987)

[7] Brenner, S., Butler, M. C. R., and King, A. D.: Periodic algebras which are almost Koszul. Algebr. Represent. Theory 5 331-367(2002)

[8] Crawley-Boevey, W., Etingof, P., Ginzburg, V.: Noncommutative Geometry and Quiver algebras, Adv. Math. 209(1) 274C336(2007)

[9] Dlab, V., Ringel, C. M.: The preprojective algebra of a modulated graph, In Representation theory II 216-231, Lecture Notes in Math 832, Springer, Berlin(1980)

[10] Geiss, Ch., Leclerc, B., Schröer, J.: Rigid modules over preprojective algebras, Invent. Math. vol 165(3) 589-632(2006)

[11] Guo, J. Y.: Translation algebras and applications, J. Alg. 255 1-21(2002)

[12] Guo, J. Y.: On McKay quivers and covering spaces. (in Chinese). Sci. Sin. Math. 41(5) 393-402(2011)

[13] Guo, J. Y.: Coverings and Truncations of Graded Self-injective Algebras, J. Alg. 355(1) 9-34(2012)

[14] Guo J Y. McKay quivers and absolutely n-complete algebras. Sci. China. Math. 56(8) 1607-1618(2013)

[15] Guo, J. Y.: On n-translation algebras. Preprint, arXiv:1406.6136

[16] Iyama, O.: Higher-dimensional Auslander-Reiten theory on maximal orthogonal subcategories, Adv. Math. 2007 210(1) 22-50(2007)

[17] Iyama, O.: Auslander correspondence, Adv. Math. 210(1) 51-82 (2007)

[18] Iyama, O.: Cluster tilting for higher Auslander algebras, Adv. Math. 226(1) 1C61(2011)

[19] Keller, B.: Cluster algebras, quiver representations and triangulated categories, In: Holm, T., Jorgensen, P., Rouquier R.:(eds.) Triangulated categories, pp. 76C160. London Math. Soc. Lecture Note Ser., vol. 375, Cambridge Univ. Press, Cambridge (2010)

[20] Reiten, I.: Cluster categories, In: Proceedings of the International Congress of Mathematicians. Volume I, 558C594, Hindustan Book Agency, New Delhi (2010)

[21] Ringel, C. M.: *Tame algebra and integral quadratic forms.* Lecture Note in Mathematics no.1099, Springer Verlag Berlin (1994)

Jin Yun Guo, Deren Luo, Department of Mathematics and Key Laboratory of HPCSIP (Ministry of Education of China), Hunan Normal University, Changsha, CHINA

E-mail address: `gjy@hunnu.edu.cn`